

## Review article

## Evidence for effective interventions to reduce mental health-related stigma and discrimination in the medium and long term: systematic review

N. Mehta,\* S. Clement,\* E. Marcus, A.-C. Stona, N. Bezborodovs, S. Evans-Lacko, J. Palacios, M. Docherty, E. Barley, D. Rose, M. Koschorke, R. Shidhaye, C. Henderson and G. Thornicroft

**Background**

Most research on interventions to counter stigma and discrimination has focused on short-term outcomes and has been conducted in high-income settings.

**Aims**

To synthesise what is known globally about effective interventions to reduce mental illness-based stigma and discrimination, in relation first to effectiveness in the medium and long term (minimum 4 weeks), and second to interventions in low- and middle-income countries (LMICs).

**Method**

We searched six databases from 1980 to 2013 and conducted a multi-language Google search for quantitative studies addressing the research questions. Effect sizes were calculated from eligible studies where possible, and narrative syntheses conducted. Subgroup analysis compared interventions with and without social contact.

**Results**

Eighty studies ( $n=422\,653$ ) were included in the review. For studies with medium or long-term follow-up (72, of which 21 had calculable effect sizes) median standardised mean differences were 0.54 for knowledge and  $-0.26$  for

stigmatising attitudes. Those containing social contact (direct or indirect) were not more effective than those without. The 11 LMIC studies were all from middle-income countries. Effect sizes were rarely calculable for behavioural outcomes or in LMIC studies.

**Conclusions**

There is modest evidence for the effectiveness of anti-stigma interventions beyond 4 weeks follow-up in terms of increasing knowledge and reducing stigmatising attitudes. Evidence does not support the view that social contact is the more effective type of intervention for improving attitudes in the medium to long term. Methodologically strong research is needed on which to base decisions on investment in stigma-reducing interventions.

**Declaration of interest**

None.

**Copyright and usage**

© The Royal College of Psychiatrists 2015. This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) licence.

Since Goffman's seminal work on stigma,<sup>1</sup> research in this field has steadily grown,<sup>2</sup> although most work consists of surveys among the general public about attitudes towards people with mental illness,<sup>3–6</sup> and much less is known about effective interventions to reduce stigma,<sup>6</sup> or about stigma in low- and middle-income countries (LMICs).<sup>7–10</sup> To better understand the evidence base on interventions to reduce mental illness-related stigma and discrimination, we identified eight existing systematic reviews on this topic.<sup>11–18</sup> The reviews varied widely in their methods and foci. There was considerable methodological and clinical heterogeneity in the included studies, and consequently meta-analysis was only undertaken in one review,<sup>11</sup> and for small subgroups in two others.<sup>12,13</sup> Four reviews presented data or commented on the overall pattern of effect sizes,<sup>11–14</sup> and in each of these the interventions had small to moderate effects, using Cohen's interpretation.<sup>19</sup> There was clearest consensus that the interventions containing social contact and first-person narratives were more effective than others.<sup>11,13,15,16</sup> Two of the reviews explored moderators of effects to understand which types of contact work best,<sup>11,13</sup> but there is a need for more research in this area. Two reviews indicated that some interventions have the potential to worsen stigma.<sup>13,17</sup> Most of the reviews were critical of the methodological quality of the included studies,<sup>12–15,18</sup> commenting in particular on the need for more randomised controlled trials (RCTs) and robust methods generally; the use

of unvalidated measures; and the relative lack of follow-up beyond the immediate post-intervention period. Other study limitations noted were the use of convenience samples,<sup>13,15,17</sup> small sample sizes,<sup>14</sup> or inappropriate outcome measures.<sup>14,15</sup> Some reviews highlighted the poor quality of the interventions, which were sometimes delivered without training, manualisation or fidelity checks,<sup>11</sup> and interventions often lacked a theoretical underpinning and developmental research.<sup>13,14</sup> In all except one review, which was restricted to studies in Iran,<sup>12</sup> interventions taking place in LMICs were a small minority or did not feature. From this scoping of existing systematic reviews we concluded that there was a need for a further systematic review to synthesise the evidence on two key issues: effectiveness in the longer term and in LMIC contexts. Consequently this systematic review aimed to assess the effectiveness of interventions (of any type with any target population), compared with inactive or baseline comparators, in reducing mental health-related stigma (knowledge, attitudes and behaviour) using any quantitative study design, addressing specifically the evidence for medium- and long-term effectiveness (research question 1) and the effectiveness of interventions in LMICs (research question 2).

**Method**

Studies were included if they described any type of intervention with a stated aim of changing mental health-related stigma or with an implied aim of changing stigma as indicated by the inclusion of

\*Joint first authors.

at least one of the following core stigma-related outcomes: stigma (any), prejudice (attitudes and related outcomes), discrimination, internalised/self-stigma or public mental health awareness/literacy. Intervention studies were included if they related to functional mental illnesses; interventions solely about, or delivered to, populations with dementia, substance misuse, intellectual disabilities or developmental disorders were excluded from this review. We included all quantitative study designs, including RCTs, controlled and uncontrolled pre–post studies, crossover studies, cohort studies and longitudinal panel studies. Studies with more than one intervention group were included. To be eligible, studies needed to report a comparison with a control group (including treatment as usual, best available current treatment or an active control, to control for non-specific effects of the intervention) or a baseline comparator. Studies needed to include at least one stigma outcome which we categorised as related to knowledge, attitudes (prejudice, self-stigma, self-esteem) or behaviour (discrimination, stigma-coping). To be eligible studies also had to address one of our two research questions: to have at least one follow-up point at least 4 weeks after the intervention was completed (to reflect the importance of medium- and longer-term outcomes relevant to stigma, as this is often described by people with mental illness as a long-term challenge); or for the intervention to be carried out in an LMIC setting. Eligibility criteria are shown in the Appendix.

### Information sources and search strategy

We identified studies by searching electronic databases, hand-checking reference lists of reviews and consulting with experts in the working group with knowledge of papers in press. We searched the following databases between 25 January 2013 and 8 February 2013: Medline, PsycINFO, the Cochrane Library, the Cumulative Index to Nursing and Allied Health Literature (CINAHL), the Social Science Citation Index (SSCI) and Global Health. In addition we conducted a Google advanced search focusing on LMICs (see Fig. 1 for details). The Google search was warranted in the light of the limited amount of stigma research in LMICs, but was precluded for our first research question as research from high-income countries is more likely to be found through a standard systematic review search. A search strategy was developed by consensus among authors (N.M., S.C., E.B. and M.D.) using both MeSH and text word searching. We searched using the format ‘Stigma’ OR ‘Discrimination’ OR synonyms AND ‘mental health’ OR ‘mental disorders’ OR synonyms AND ‘Intervention Studies’ OR synonyms. The full Medline search strategy is shown in online Table DS1. The search was restricted to results between 1980 and 2013 and studies on human beings, but was not limited by language. The decision to start the search at 1980 was a pragmatic one based on our examination of the existing reviews which revealed that the vast majority of stigma intervention research commenced after 1980. Relevant non-English language papers were read by fluent native language speakers in French and Spanish according to the linguistic skills available to members of the review team. Potentially relevant papers in many important languages, including Chinese, were therefore excluded from the review. Systematic and non-systematic reviews were identified during the search and the reference lists of these studies were hand-checked.

### Study selection and data collection

All identified titles and abstracts were screened by two researchers. Because of the large number of search hits, two researchers screened 5% of abstracts together. As good agreement (> 95%) was achieved, the remainder were divided between the two

researchers and study selection conducted by one researcher for each half. Where the researcher was unclear as to whether a paper should be included, the paper was discussed in consensus meetings. Two review authors extracted data from included studies for all parts of the systematic review, with queries resolved by discussion and consensus.

### Statistical analysis

Outcomes for the studies included were reported using both scales and individual items, although for the effect size calculations were restricted to scale data for knowledge and attitudes. We classified all reported stigma outcomes into the categories of ‘knowledge’, ‘attitudes’ or ‘behaviour’. Differences between intervention group and control group at follow-up were our main focus for the quantitative review. Effect sizes, standardised mean differences (SMDs) and 95% confidence intervals were calculated for studies where there were sufficient data to calculate this using the Campbell Collaboration effect size calculator.<sup>20</sup> We had planned to calculate odds ratios for dichotomous outcomes but found no study for which this was calculable. Negative SMDs indicate a reduction in stigma (benefit), i.e. an improvement in knowledge outcomes or a reduction in either negative attitudes or discriminatory behaviour in the intervention group. Where more than one outcome was reported within a category, the median effect size was presented.<sup>21</sup> In the online tables we present data on the number of outcomes with statistically significant changes in outcome and the direction of effect to complement the effect size data of outcomes.<sup>21</sup> These also provide some information about all included studies and at least some information on effectiveness for studies that reported insufficient data to calculate effect sizes. Owing to the considerable heterogeneity of the interventions, measures and participants in the included studies, it was not possible to conduct meta-analyses or to use conventional analytical methods to control for heterogeneity. As some studies had more than one intervention, this analysis was carried out at the intervention level with the number of participants in the control group split between the interventions, to control for unit of analysis error.<sup>22</sup>

We conducted two subgroup analyses on type of intervention by calculating, presenting and comparing median effect sizes attitude outcomes for each subgroup. The first analysis compared direct, indirect or no social contact, and the second compared target groups. We undertook similar sensitivity analyses to explore the possible effects of study design and risk of bias. First, we compared RCT evidence with non-RCT evidence, and second, within RCTs we compared the third of studies with the least risk of bias (see below) with the remainder.

### Assessment of study quality and risk of bias

A quality assessment and profile of risk of bias within studies were carried out individually for all included studies. Level of RCT evidence was rated by two authors using the Cochrane risk of bias tool.<sup>23</sup> The third of RCTs with the lowest risk of bias are identified with an asterisk in the data extraction tables. To assess bias in non-randomised studies two researchers conducted quality appraisals using risk of bias criteria for non-randomised studies,<sup>23</sup> suitable to the wide range of study designs included. When a decision about the risk of bias could not be made, it was resolved through discussion with a third author. In addition, for each study we indicated whether at least one outcome measure was validated, whether it was previously published, developed by the author or if items were used. Scales were marked as having evidence of psychometric adequacy providing they met one or more of the following

criteria: the authors reported a Cronbach's  $\alpha$  of 0.7 or greater, the authors referenced the measure as being reliable or valid, or there was some evidence of validity or reliability as judged by the review team.

## Results

A total of 80 quantitative studies (422 653 participants) were identified for inclusion in the review, 72 addressing research question 1 (long-term effectiveness) and 11 addressing research question 2 (setting), of which 3 studies addressed both questions (Fig. 1). The database search provided 27 876 citations. After a review of the abstracts 26 563 papers were excluded as they were clearly irrelevant or did not meet the inclusion criteria. The reference lists of 17 reviews were hand-checked and 49 further papers identified. Seven papers in press known to the authors were included. After removal of 330 duplicates the full text of the remaining 1061 potentially relevant papers was sought. Of these, 21 papers were unobtainable and 843 papers did not meet the inclusion criteria. Of the remaining papers 17 did not contain enough relevant data to extract. A full reference list is given in online Table DS2 and study characteristics are listed in online Tables DS3–5. Online Tables DS6 and DS7 give risk of bias and quality ratings for RCTs and other studies respectively.

## Medium- and long-term follow-up

### Study characteristics

Most of the studies addressing medium- or long-term outcomes took place in high-income countries (93%), were aimed at school or university students (37%) and used interventions comprising mental health education and literacy or mental health information (43%). About a quarter (28%) of the studies included were RCTs, 52% consisted of pre–post studies with or without a control group and 21% were longitudinal panel or cohort studies. Most studies (69%) had a final follow-up assessment 1–6 months after the intervention had ended, whereas 21% had a longer follow-up (1–10 years post-intervention). Tables DS3 and DS4 show details of study characteristics.

### Evidence

There were 72 quantitative studies with at least 4 weeks of follow-up, which included 81 interventions with 42 653 participants. It was possible to calculate effect sizes and confidence intervals for 21 of these studies (23 interventions). These studies and their effect sizes are shown in Table 1. Findings based on statistical significance for all included studies are shown in Tables




Fig. 1 Selection of papers and sources included in the review. LMIC, low- and middle-income country; SSCI, Social Science Citation Index.

DS3 (RCT, controlled and uncontrolled pre–post studies) and DS4 (longitudinal panel study or cohort design). For knowledge outcomes the median effect size was 0.54 indicating a medium effect in increasing knowledge.<sup>19</sup> For attitude outcomes SMDs ranged from 0.05 to  $-1.22$  with a median effect size of  $-0.26$ , indicating a small reduction in stigmatising attitudes. For behavioural outcomes SMDs were calculated in one intervention which showed a small (SMD = 0.22) effect in reducing stigmatising behaviour. Inspection of the pattern of significance findings for scales for all the included studies with medium- or long-term follow-up indicated that there were similar numbers of significant and non-significant findings indicating an increase in knowledge (26 *v.* 22). Similar numbers were also found for attitude scales (63 non-significant findings *v.* 52 significant in the direction of stigma reduction). Five scales had significant findings indicating an increase in stigma. For behavioural outcomes measured with scales, non-significant findings outnumbered significant ones indicating a reduction in discriminatory behaviour (12 *v.* 2) and this was also the case for behavioural outcomes measured at the item level (38 *v.* 19).

Our subgroup analysis of type of intervention found that interventions containing direct social contact had a smaller median effect size for stigmatising attitudes ( $-0.17$ ) than those with indirect social contact ( $-0.32$ ) or no social contact ( $-0.33$ ). There were enough interventions with effect sizes to make comparisons of median effect sizes by three types of target group, and we found that interventions targeted at health professionals had a somewhat higher median effect size ( $-0.41$ ) than those targeting school pupils ( $-0.21$ ) or university students ( $-0.13$ ).

#### Risk of bias

Across all RCTs there was a low risk of bias for 50% of the criteria and an unclear or high risk of bias in the other 50%. Only five trials met 70% or more of the criteria. Nine trials met between 40% and 60% of the criteria and five only met 15–30%. In light of the nature of anti-stigma interventions it was not possible to mask participants and personnel to allocation, with the exception of one trial which was internet-based and thus easier to conceal.<sup>24</sup> Of the 19 trials, 17 used at least one validated scale to measure outcomes, whereas 2 used non-validated scales that had been used in previously published papers. There were 53 non-randomised studies, 30 of which did not have a control group. Among studies with a control group, 6 were deemed to have a low risk of selection bias with regard to the comparability between the intervention and control groups. In 26 studies there was a high risk of attrition bias, where more than 20% of the sample were lost to follow-up and no intention to treat analysis was carried out. Possible confounders were considered and controlled for in only 28% of studies. As with the RCTs, masking of participants and personnel was not possible owing to the type of intervention. Among non-randomised studies, 24 had at least one validated outcome measure, 2 had at least one that was previously published, 4 had one that was specifically developed for the study with no psychometric testing reported, and 23 used items only. Details of risk of bias in individual studies are given in Tables DS5 and DS6. The median effect size for RCTs was lower than for non-randomised controlled studies ( $-0.17$  *v.*  $-0.37$ ). Within RCTs the third with the least risk of bias had a higher effect size ( $-0.30$ ) than the remainder ( $-0.09$ ).

#### Evidence from LMICs

There were 11 studies (1967 participants) from LMIC settings, 8 with less than a 4-week follow-up and 3 with longer follow-up. Study characteristics and statistical significance findings for these

are shown in Table DS5. Eight of these were from upper middle-income countries and three were from lower middle-income countries. There was no study meeting our criteria from a low-income country. Six studies were aimed at school and university students, two at caregivers of people with schizophrenia, and three at healthcare professionals. Three studies used an RCT design, one of which was a cluster randomised trial analysed within groups, two were controlled studies and six were uncontrolled pre–post studies. Within the 11 studies included there were 16 intervention arms, with 5 measuring knowledge outcomes and 14 measuring attitude outcomes. None of the studies had behavioural outcomes. Sufficient data to calculate an effect size were reported in only one of the studies;<sup>25</sup> in this study – a psychoeducation programme for caregivers of patients with schizophrenia in Chile – the SMD for stigmatising attitudes was  $-2.11$  (95% CI  $-2.87$  to  $-1.34$ ), indicating a large effect. Inspection of the statistical significance of the knowledge scale findings for all studies revealed that both studies with such outcomes found no evidence of change; however, there were findings indicating a significant reduction in stigmatising attitudes for 11 of the 12 attitude scale outcomes assessed in these studies (Table DS5).

These results should be interpreted with caution. In seven of the studies, follow-up assessments were undertaken immediately after the intervention (in one study this was done 1 week after the intervention had ended). There were also issues regarding bias: owing to a lack of information in the papers it was generally difficult to gauge the extent of risk of bias. For the three RCTs, in 52% of criteria the risk of bias was unclear. Where information was provided, a high risk of bias was found in 19% of criteria across the RCTs, whereas in 29% of criteria the risk was low. This was most common for the incomplete outcome data and selective outcome reporting criteria. For the non-randomised studies, risk of bias varied across criteria, with 33% classified as high and 33% as low, and for 33% the degree of risk was unclear.

#### Behavioural outcomes

Among the 15 studies that did report behavioural outcomes, 7 assessed contact with someone with a mental health problem, 4 measured perceived discrimination and coping strategies in participants who had a mental health problem, 2 measured changes in school and workplace policies regarding mental health,<sup>26,27</sup> 2 measured experienced discrimination reported by people with mental health problems,<sup>28,29</sup> and only 1 measured actual discriminatory behaviour by participants in the general population.<sup>30</sup>

## Discussion

Our synthesis of 72 studies with follow-up beyond 4 weeks revealed that, at this follow-up, interventions aimed at reducing mental health-related stigma typically had a medium-sized effect on knowledge outcomes and a small effect on attitudinal outcomes, although for both types of outcome statistically non-significant findings were as common as significant ones. There were insufficient data on behavioural outcomes to draw any conclusions on the medium- or long-term effectiveness of interventions to reduce discrimination. This is the first systematic review to synthesise evidence on medium- and long-term effectiveness, which is striking given that stigma is often experienced by people with mental illness as a long-term difficulty. Although a number of systematic reviews indicated that social contact interventions were particularly effective,<sup>11,13,15,16</sup> the majority of studies in these reviews had only short-term follow-up. Our review, restricted to studies with medium- and

**Table 1** Evidence for medium- and long-term effectiveness of interventions to reduce mental health-related stigma

Study <sup>a</sup>	Design <sup>b</sup>	n <sup>c</sup>	Intervention	Time to follow-up <sup>d</sup>	Evidence for effectiveness SMD (95% CI) <sup>e</sup>		
					Knowledge	Attitudes	Behaviour
Targeted at the armed forces							
Seal <i>et al</i> (2012) <sup>45</sup>	RCT	73	Motivational interviewing	8 weeks		0.04 (−0.07 to 0.86)	
Gould <i>et al</i> (2007) <sup>46</sup>	Controlled	124	Training programme to provide support, education and modify attitudes about PTSD	1 month		0.42(r) (0.00 to 0.85)	
Targeted at school students							
Campbell <i>et al</i> (2011) <sup>47</sup>	RCT	92	Mental health workshop including education and direct contact	10 weeks		0.05 (−0.39 to 0.49)	
Pinto-Foltz <i>et al</i> (2011) <sup>48</sup>	RCT	156	Direct contact with service users who were in sustained recovery from mental illness	8 weeks	0.29 (−0.05 to 0.63)	−0.17 (−0.50 to 0.17)	
Esters <i>et al</i> (1998) <sup>49</sup>	Controlled	40	Mental health education about stigma and help-seeking	12 weeks		<b>−0.45</b> <b>(−1.08 to −0.18)</b>	
O’Kearney <i>et al</i> (2006) <sup>50</sup>	Controlled	59	Internet programme aiming to help people identify, overcome and cope with depression	16 weeks		−0.25 (−0.83 to 0.34)	
O’Kearney <i>et al</i> (2009) <sup>51</sup>	Controlled	157	Internet programme aiming to help people identify, overcome and cope with depression	20 weeks	−0.14 (−0.45 to 0.18)	<b>−0.17</b> <b>(−0.49 to 0.15)</b>	
Ventieri <i>et al</i> (2011) <sup>52</sup>	Controlled	195	Mental health education, with role play and activities	4 months	<b>0.51</b> <b>(0.21 to 0.80)</b>	<b>−0.33</b> <b>(−0.62 to −0.03)</b>	
Targeted at university students							
Gonzales <i>et al</i> (2002) <sup>53</sup>	RCT	167	Mental health education about stigma	4 weeks		−0.07 (−0.52 to 0.38)	
Sharp <i>et al</i> (2006) <sup>54</sup>	RCT	123	Mental health education	1 month		−0.09 (−0.47 to 0.29)	
Faigin & Stein (2008) <sup>55</sup>	Controlled	204	A play by actors with history of severe mental illness addressing their experiences and stigma	1 month		−0.13 (−0.47 to 0.20)	
Faigin & Stein (2008) <sup>55</sup> (2nd arm)	Controlled	222	A video-recorded version of the play described above	1 month		−0.37 (−0.69 to −0.05)	
O’Reilly <i>et al</i> (2011) <sup>56</sup>	Controlled	272	Mental health first aid training for pharmacy students	6 weeks		−0.61 (−0.92 to −0.31)	
Targeted at healthcare professionals							
Blair Irvine <i>et al</i> (2012) <sup>57</sup>	RCT	172	Internet courses with behavioural skills and knowledge training for long-term care staff	1 month	<b>0.56</b> <b>(0.25 to 0.86)</b>	<b>−0.17</b> <b>(−0.47 to 0.13)</b>	
Patterson <i>et al</i> (2007) <sup>58</sup>	Controlled <sup>f</sup>	91	Educational intervention about self-harm behaviour for nurses	18 months		−1.22 (−1.86 to −0.58)	
Treloar (2009) <sup>59</sup>	Controlled <sup>f</sup>	90	Educational programme about self-harm using psychoanalytic aetiology framework	6 months		−0.35 (−1.06 to 0.37)	
Treloar (2009) <sup>59</sup> (2nd arm)	Controlled <sup>f</sup>	91	Educational programme about self-harm using CBT aetiology framework	6 months		−0.47 (−1.23 to 0.29)	
Targeted at the general public							
Jorm <i>et al</i> (2004) <sup>60</sup>	RCT*	753	Mental health first aid course	4 months	<b>11.77</b> <b>(5.98 to 17.56)</b>	<b>−0.26 (−0.49 to −0.03)</b>	0.22(r) (−0.18 to 0.63)
Targeted at people with mental health problems							
Fung <i>et al</i> (2011) <sup>61</sup>	RCT*	66	Self-stigma reduction programme	6 months		0.34 (−0.82 to 0.15)	
Gumley <i>et al</i> (2006) <sup>62</sup>	RCT	144	CBT targeting negative beliefs about self and illness	12 months		−0.12 (−0.45 to 0.21)	
Targeted at other groups							
Gulliver <i>et al</i> (2012) <sup>24</sup>	RCT*	59	Mental health literacy and destigmatisation intervention for elite athletes	3 months	0.76 (−0.17 to 1.68)	0.50(r) (0.41 to 1.41)	
Kitchener & Jorm (2004) <sup>27</sup>	RCT*	301	Mental health first aid course for employees	5 months	0.07 (−0.16 to 0.30)	−0.17 (−0.40 to 0.05)	
Jorm <i>et al</i> (2010) <sup>63</sup>	RCT*	327	Youth mental health first aid course for teachers	6 months	0.67 (0.18 to 0.65)		

CBT, cognitive-behavioural therapy; PTSD, post-traumatic stress disorder; RCT, randomised controlled trial; SMD, standardised mean difference.

a. Studies with sufficient data to calculate effect sizes.

b. Designs include RCTs in the top tercile for quality, i.e. highest numbers of Cochrane risk of bias items rated as low (RCT\*); RCTs in the lower two terciles for quality (RCT) (see online Table DS5 for details); pre-post studies with a control group (Controlled).

c. Number of participants in the intervention and control groups.

d. Time to final follow-up results.

e. An SMD <0 indicates a reduction in knowledge, stigmatising attitudes or stigmatising behaviours unless the data are such that this can only be calculated to show the reverse effect, in which case this is marked (r). Bold type indicates confidence intervals that do not cross zero.

f. Hedges’ g used by study authors instead of Cohen’s d owing to small sample sizes.

longer-term outcomes, did not support the superiority of social contact interventions as we had expected. As it is vital that stigma reduction is sustained in the longer term, the effectiveness of such social contact interventions clearly warrants further research.

Study quality was variable, and indeed study design and quality did appear to affect median effect sizes, although these subgroup and sensitivity analysis findings should be interpreted with caution owing to the heterogeneity of the studies. Overall, where we did identify positive changes from the interventions, the magnitude of the effects was generally rather modest. It is also clear that there is therefore a lack of research on actual discriminatory behaviour within the stigma research field.

For our second research question regarding LMICs, we found comparatively few studies from middle-income countries and none from low-income countries. A large effect size was found for the one LMIC study for which there were sufficient data to calculate the effect size and the majority of attitude scale outcomes indicated significant improvements in attitudes, although such findings must be treated with considerable caution. There is a clear need for more stigma reduction studies, particularly from low-income countries.

Our results regarding service user social contact are consistent with those of Griffiths *et al.*,<sup>31</sup> who recently published a meta-analysis of RCTs of interventions intended to reduce stigma. Analysing data from 26 trials they found that interventions targeting personal stigma or social distance yielded small but significant reductions in stigma across all mental disorders. Further, they reported that educational interventions were effective in reducing personal stigma, as were interventions incorporating service user contact. This study also considered internet use and self-stigma and found that internet programmes were at least as effective in reducing personal stigma as face-to-face delivery (see also Clement *et al.*).<sup>32</sup> They found no evidence that stigma interventions were effective in reducing self-stigma. In our review, although social contact appears to be the most strongly evidence-based type of intervention to reduce stigma when measured by immediate post-intervention outcomes, there is not at present evidence to show that such immediate benefits persist in the longer term.

### Limitations of the study

This review has a number of limitations. In conducting a comprehensive overview of all relevant literature we have identified considerable heterogeneity among participant groups, interventions and outcomes. For example, we identified 55 different scales used for the 136 outcomes measured. Study quality also varied considerably. We were able to include studies in some non-English languages, but it is possible that we missed important projects published in other languages, for example potentially important studies not available at all in English, or studies for which only abstracts were available in English, and which we were not able to assess fully (see, for example, Shi *et al.*).<sup>33</sup> The temporal limitation of the search start date being 1980 will have resulted in the review missing studies before that date. We also need to acknowledge the possibility of publication bias, for example that intervention studies showing no difference might be published less often than those that do identify a clear benefit. Further, the risk of bias results given above, with half of all studies having a high or unknown risk of bias, mean that considerable caution needs to be exercised in interpreting these findings. It is also notable that relatively few of the interventions assessed followed published, manualised procedures or including any rating of treatment fidelity. It should also be appreciated that although a narrative review may be able to disaggregate the nature of the interventions, and the specific target groups, into a greater number of specific subtypes, the numbers

of studies in each of these categories would be small, and that this would give a greater descriptive richness at the expense of the wider generalisability of the findings. The systematic review method used here does not allow this narrower focus.

### Challenges in the measurement of stigma

The assessment and validation of instruments to measure stigma and discrimination against people with mental illness has been under way since the 1960s. Although early measures such as the Opinions About Mental Illness and the Community Attitudes to Mental Illness scales are still used in some studies,<sup>34,35</sup> there have been many developments in the breadth and quantity of measures to assess stigma in recent years. These include a trend to incorporate multiple outcomes or domains, for example knowledge and behaviour as well as attitudes; techniques to control for social desirability bias such as implicit measures; research on coping or 'stigma resilience'; and assessments among multiply stigmatised groups, such as people from ethnic minorities with mental illness. Despite these developments there are still substantial gaps in what can be assessed using available measures, including a lack of behavioural and structural indicators. We have seen in this review that behaviour is under-represented in stigma intervention outcomes, for example changes in behaviour of others rated by patients or service users, or directly observed discrimination-related outcomes. There is a further gap in terms of important subgroups. For example, Link *et al* noted that children and adolescents were represented in only 3.7% of stigma studies.<sup>36</sup> More specific and tailored measures might facilitate inclusion of specific subpopulations in stigma research, such as those already affected by discrimination on the grounds of (for example) ethnicity. Additionally, studies that include measures validated in LMICs are rare, and only a few include any intervention component developed specifically in such countries. Future efforts should therefore address these gaps, because measurement and evaluation are critical to understanding the underlying mechanisms and effectiveness of anti-stigma interventions. A further challenge is to stop the use of unvalidated measures and item level analyses, while retaining enough flexibility to promote conceptual, contextual and theoretical relevance.

### Gaps in the evidence base

This review has highlighted clear gaps in the field of anti-stigma interventions and research methods and a need for the harmonisation of outcomes in this field of research. These include the paucity of evidence on discrimination outcomes, or on reducing negative behaviours or increasing positive behaviours towards people with mental illness,<sup>37</sup> and the lack of studies of specific target groups such as employers or family members, despite service users commonly reporting experiencing discrimination from both of these groups.<sup>38</sup> There is an important need to assess whether benefits identified in the short term are maintained in the longer term, and if any booster interventions are needed to achieve sustainability. This review has also shown a relatively narrow focus of work to date: either on the general population (in attitude surveys) or on students within settings accessible to researchers (e.g. universities and colleges).<sup>16</sup> From a global health viewpoint there is a distinct lack of interventional research in LMICs, despite emerging evidence of the scale and severity of the challenges posed by stigma and discrimination, and despite the fact that 85% of the world's population live in such countries.<sup>39,40</sup> Finally, there is a need for more studies using high-quality research designs. Only a third of studies included in this paper used an RCT or other robust study design, and many of these had a high risk of bias.

## Future research

Knowledge in this field is generally from small studies of poor methodological quality, using inconsistent outcomes scales, and in particular few strong RCTs or interrupted time series studies have been carried out to test interventions intended to reduce stigma and discrimination. Our summary of previous systematic reviews does tend to support the view that social contact is the more effective type of intervention known to reduce stigma, at least in the short term.<sup>41</sup> We do not yet have even weak consistent evidence to support interventions for target groups identified as priorities by service user groups, such as family members, and only an embryonic evidence base concerning how to address stigma in healthcare staff.<sup>42</sup> Indeed, this degree of evidential neglect could itself be seen as a manifestation of structural discrimination. Given the magnitude of the challenges posed by stigma and discrimination, it is clear that there needs to be a commensurate concerted effort to fund methodologically strong research to provide robust evidence to support policy decisions on investment and interventions. Such a wider policy framework is now emerging.<sup>43</sup> The World Health Organization Mental Health Action Plan, ratified by the World Health Assembly in May 2013, states as its vision:

'A world in which mental health is valued, promoted and protected, mental disorders are prevented and persons affected by these disorders are able to exercise the full range of human rights and to access high quality, culturally-appropriate health and social care in a timely way to promote recovery, in order to attain the highest possible level of health and participate fully in society and at work, free from stigmatization and discrimination.'<sup>44</sup>

Specifically, paragraph 75 of the Action Plan indicates a need to prioritise:

'Mental health promotion and prevention: provide technical support to countries on the selection, formulation and implementation of evidence-based and cost-effective best practices for promoting mental health, preventing mental disorders, reducing stigmatization and discrimination, and promoting human rights across the lifespan.'<sup>44</sup>

This review indicates that an early necessity is to conduct more high-quality research to allow this policy priority to be firmly evidence-based, especially within LMICs.

**Nisha Mehta**, MBBS, **Sarah Clement**, PhD, **Elena Marcus**, MSc, **Anne-Claire Stona**, BSc, **Nikita Bezborodovs**, MD, **Sara Evans-Lacko**, PhD, **Jorge Palacios**, PhD, **Mary Docherty**, MRCPsych, **Elizabeth Barley**, PhD, **Diana Rose**, PhD, **Mirja Koschorke**, PhD, Health Service and Population Research Department, King's College London, Institute of Psychiatry, Psychology and Neuroscience, London, UK; **Rahul Shidhaye**, PhD, Centre for Mental Health, Public Health Foundation of India, Delhi, India; **Claire Henderson**, PhD, **Graham Thornicroft**, PhD, Health Service and Population Research Department, King's College London, Institute of Psychiatry, Psychology and Neuroscience: London, UK

**Correspondence:** Professor Graham Thornicroft, Health Service and Population Research Department, Box PO29, Institute of Psychiatry, Psychology and Neuroscience, De Crespigny Park, London SE5 8AF, UK. Email: graham.thornicroft@kcl.ac.uk

First received 20 Sep 2013, final revision 4 Jan 2015, accepted 27 Feb 2015

## Funding

This publication is independent research funded by the National Institute for Health Research (NIHR) under its Programme Grants for Applied Research scheme (Improving Mental Health Outcomes by Reducing Stigma and Discrimination: RP-PG-0606-1053) (authors G.T., S.C., C.H., D.R.). This research was supported by the NIHR Collaboration for Leadership in Applied Health Research and Care South London at King's College London Foundation Trust. G.T. is supported by the European Union Seventh Framework Programme (FP7/2007-2013) Emerald project. This paper is an output of the Programme for Improving Mental health care (PRIME). The material has been funded by aid from the UK Government; however, the views expressed do not necessarily reflect the UK Government's official policies. The research supported by the NIHR Collaboration for Leadership in Applied Health Research and Care South London at King's College London Foundation Trust. G.T. is also funded in relation to the NIHR Specialist Mental Health Biomedical Research Centre at the Institute of Psychiatry, Psychology and Neuroscience, King's College London and the South London and Maudsley National Health Service (NHS) Foundation Trust. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health. M.K. was supported by a Wellcome Trust Clinical PhD Fellowship during her involvement in the review. N.B. was funded by an Erasmus grant from the European Commission. The work leading to these results has received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement 305968.

## Acknowledgements

We acknowledge our gratitude to the following colleagues for their contributions to this paper: Luigi de Benedictis and Dmitry Krupchanka (grey literature search); Nikhil Gupta (risk of bias assessments and referencing); Henry Lishi Li (extraction of one Chinese paper); and Shinsuke Koike (helped to extract one Japanese paper).

## Appendix

### Eligibility criteria for study inclusion

#### Participants

Any, except target populations that solely comprised people with dementia, substance misuse, intellectual disabilities or developmental disorders.

#### Setting

Any.

#### Intervention

Any intervention with a stated aim of changing mental health-related stigma, or with an implied aim of changing stigma as indicated by the inclusion of at least one of the following stigma-related outcomes: stigma (including internalised stigma), prejudice (attitudes and related outcomes), discrimination, or public mental health awareness/mental health literacy. Interventions relating to functional mental illnesses were included, those solely about dementia, substance misuse, learning disabilities or developmental disorders were excluded.

#### Comparison

Inactive or baseline comparator.

#### Outcomes

Outcomes comprising:

- knowledge
- attitudes (prejudice/self-attitudes)
- behaviour (discrimination/stigma-coping)
- follow-up at least 4 weeks after the intervention was completed (research question 1) or any (research question 2).

#### Study design

Any quantitative design.

## References

- 1 Goffmann E. *Stigma: Notes on the Management of Spoiled Identity*. Prentice Hall, 1963.
- 2 Weiss MG, Ramakrishna J, Somma D. Health-related stigma: rethinking concepts and interventions. *Psychol Health Med* 2006; **11**: 277–87.
- 3 Sartorius N, Schulze H. *Reducing the Stigma of Mental Illness*. Cambridge University Press, 2005.
- 4 Rabkin J. Public attitudes toward mental illness: a review of the literature. *Schizophr Bull* 1974; **10**: 9–33.
- 5 Link BG, Phelan JC, Bresnahan M, Stueve A, Pescosolido BA. Public conceptions of mental illness: labels, causes, dangerousness, and social distance. *Am J Public Health* 1999; **89**: 1328–33.
- 6 Thornicroft G. *Shunned: Discrimination Against People with Mental Illness*. Oxford University Press, 2006.
- 7 Yang LH, Cho SH, Kleinman A. Stigma of mental illness. In *Mental and Neurological Public Health: A Global Perspective* (ed. V Patel): 484–95. Elsevier, 2010.
- 8 Yang LH, Kleinman A, Link BG, Phelan JC, Lee S, Good B. Culture and stigma: adding moral experience to stigma theory. *Soc Sci Med* 2007; **64**: 1524–35.
- 9 Raguram R, Raghu TM, Vounatsou P, Weiss MG. Schizophrenia and the cultural epidemiology of stigma in Bangalore, India. *J Nerv Ment Dis* 2004; **192**: 734–44.
- 10 Weiss MG, Jadhav S, Raguram R, Vounatsou P, Littlewood R. Psychiatric stigma across cultures: local validation in Bangalore and London. *Anthropol Med* 2001; **8**: 71–87.
- 11 Corrigan PW, Morris SB, Michaels PJ, Rafacz JD, Rusch N. Challenging the public stigma of mental illness: a meta-analysis of outcome studies. *Psychiatr Serv* 2012; **63**: 963–73.
- 12 Mansouri N, Gharaee B, Shariat SV, Bolhari J, Nooraie RY, Rahimi-Movaghar A, et al. The change in attitude and knowledge of health care personnel and general population following trainings provided during integration of mental

- health in primary health care in Iran: a systematic review. *Int J Ment Health Syst* 2009; **3**: 15.
- 13 Clement S, Lassman F, Barley E, Evans-Lacko S, Williams P, Yamaguchi S, et al. Mass media interventions for reducing mental health-related stigma. *Cochrane Database Syst Rev* 2013; **7**: CD009453.
  - 14 Mittal D, Sullivan G, Chekuri L, Allee E, Corrigan PW. Empirical studies of self-stigma reduction strategies: a critical review of the literature. *Psychiatr Serv* 2012; **63**: 974–81.
  - 15 Holzinger A, Dietrich S, Heitmann S, Angermeyer M. Evaluation of target-group oriented interventions aimed at reducing the stigma surrounding mental illness. *Psychiatr Prax* 2008; **35**: 376–86.
  - 16 Yamaguchi S, Wu SI, Biswas M, Yate M, Aoki Y, Barley EA, et al. Effects of short-term interventions to reduce mental health-related stigma in university or college students: a systematic review. *J Nerv Ment Dis* 2013; **201**: 490–503.
  - 17 Ando S, Clement S, Barley E, Thornicroft G. The simulation of hallucinations to reduce the stigma of schizophrenia: a systematic review. *Schizophr Res* 2011; **133**: 8–16.
  - 18 Schachter HM, Girardi A, Ly M, Lacroix D, Lumb AB, van Berkow J, et al. Effects of school-based interventions on mental health stigmatization: a systematic review. *Child Adolesc Psychiatr Ment Health* 2008; **2**: 18.
  - 19 Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. Erlbaum, 1988.
  - 20 Wilson DB. *Practical Meta-Analysis Effect Size Calculator*. Campbell Collaboration, 2001 ([http://www.campbellcollaboration.org/resources/effect\\_size\\_input.php](http://www.campbellcollaboration.org/resources/effect_size_input.php)).
  - 21 Grimshaw J, McAuley LM, Bero LA, Grilli R, Oxman AD, Ramsay C, et al. Systematic reviews of the effectiveness of quality improvement strategies and programmes. *Qual Saf Health Care* 2003; **12**: 298–303.
  - 22 Higgins JPT, Green S (eds). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.0.2, ch 16.5. Cochrane Collaboration, 2009.
  - 23 Higgins JP, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011; **343**: 5928.
  - 24 Gulliver A, Griffiths KM, Christensen H, Mackinnon A, Calear AL, Parsons A, et al. Internet-based interventions to promote mental health help-seeking in elite athletes: an exploratory randomized controlled trial. *J Med Internet Res* 2012; **14**: 120–37.
  - 25 Gutierrez-Maldonado J, Caqueo-Urizar A, Ferrer-Garcia M. Effects of a psychoeducational intervention program on the attitudes and health perceptions of relatives of patients with schizophrenia. *Soc Psychiatry Psychiatr Epidemiol* 2009; **44**: 343–8.
  - 26 Jorm AF, Kitchener BA, Fischer JA, Cvetkovski S. Mental health first aid training by e-learning: a randomized controlled trial. *Aust NZ J Psychiatry* 2010; **44**: 1072–81.
  - 27 Kitchener BA, Jorm AF. Mental health first aid training in a workplace setting: a randomized controlled trial. *BMC Psychiatry* 2004; **4**: 23.
  - 28 Corker E, Hamilton S, Henderson C, Weeks C, Pinfold V, Rose D, et al. Experiences of discrimination among people using mental health services in England 2008–11. *Br J Psychiatry* 2013; **202** (suppl 55): s58–63.
  - 29 Henderson C, Corker E, Lewis-Holmes E, Hamilton S, Flach C, Rose D, et al. England's time to change antistigma campaign: one-year outcomes of service user-rated experiences of discrimination. *Psychiatr Serv* 2012; **63**: 451–7.
  - 30 Evans-Lacko S, Henderson C, Thornicroft G. Public knowledge, attitudes and behaviour regarding people with mental illness in England 2009–2012. *Br J Psychiatry* 2013; **202** (suppl 55): s51–7.
  - 31 Griffiths KM, Carron-Arthur B, Parsons A, Reid R. Effectiveness of programs for reducing the stigma associated with mental disorders. A meta-analysis of randomized controlled trials. *World Psychiatry* 2014; **13**: 161–75.
  - 32 Clement S, van Nieuwenhuizen A, Kassam A, Flach C, Lazarus A, de Castro M, et al. Filmed v. live social contact interventions to reduce stigma: randomised controlled trial. *Br J Psychiatry* 2012; **201**: 57–64.
  - 33 Shi ZY, Liu SY, Nie LY. The impact of group psychological interventions on the stigma perceived of patients with depression. *J Nurs Training* 2012; **27**: 1213–1216.
  - 34 Cohen J, Struening EL. Opinions about mental illness in the personnel of two large mental hospitals. *J Abnorm Soc Psychol* 1962; **64**: 349–60.
  - 35 Taylor SM, Dear MJ, Hall GB. Attitudes toward the mentally ill and reactions to mental health facilities. *Soc Sci Med* 1979; **13**: 281–90.
  - 36 Link BG, Yang LH, Phelan JC, Collins PY. Measuring mental illness stigma. *Schizophr Bull* 2004; **30**: 511–41.
  - 37 Sharac J, McCrone P, Clement S, Thornicroft G. The economic impact of mental health stigma and discrimination: a systematic review. *Epidemiol Psychiatr Soc* 2010; **19**: 223–32.
  - 38 Henderson C, Williams P, Little K, Thornicroft G. Mental health problems in the workplace: changes in employers' knowledge, attitudes and practices in England 2006–2010. *Br J Psychiatry* 2013; **202** (suppl 55): s70–6.
  - 39 Thornicroft G, Brohan E, Rose D, Sartorius N, Leese M. Global pattern of experienced and anticipated discrimination against people with schizophrenia: a cross-sectional survey. *Lancet* 2009; **373**: 408–15.
  - 40 Lasalvia A, Van Weeghel J, Reneses B, Bacle SV, Thornicroft G, Bonetto C. Discrimination reported by people with major depressive disorder: authors' reply. *Lancet* 2013; **381**: 1181–2.
  - 41 Henderson C, Thornicroft G. Evaluation of the Time to Change programme in England 2008–2011. *Br J Psychiatry* 2013; **202** (suppl 55): s45–8.
  - 42 Jones S, Howard L, Thornicroft G. 'Diagnostic overshadowing': worse physical health care for people with mental illness. *Acta Psychiatr Scand* 2008; **118**: 169–71.
  - 43 United Nations. *Convention on the Rights of Persons with Disabilities*. UN, 2006.
  - 44 World Health Organization. *Global Mental Health Action Plan 2013–2020*. WHO, 2013.
  - 45 Seal KH, Abadjian L, McCamish N, Shi Y, Tarasovsky G, Weingardt K. A randomized controlled trial of telephone motivational interviewing to enhance mental health treatment engagement in Iraq and Afghanistan veterans. *Gen Hosp Psychiatry* 2012; **34**: 450–9.
  - 46 Gould M, Greenberg N, Hetherington J. Stigma and the military: evaluation of a PTSD psychoeducational program. *J Trauma Stress* 2007; **20**: 505–15.
  - 47 Campbell M, Shryane N, Byrne R, Morrison AP. A mental health promotion approach to reducing discrimination about psychosis in teenagers. *Psychosis* 2011; **3**: 41–51.
  - 48 Pinto-Foltz MD, Logsdon MC, Myers JA. Feasibility, acceptability, and initial efficacy of a knowledge-contact program to reduce mental illness stigma and improve mental health literacy in adolescents. *Soc Sci Med* 2011; **72**: 2011–9.
  - 49 Esters IG, Cooker PG, Ittenbach RF. Effects of a unit of instruction in mental health on rural adolescents' conceptions of mental illness and attitudes about seeking help. *Adolescence* 1998; **33**: 469–76.
  - 50 O'Kearney R, Gibson M, Christensen H, Griffiths KM. Effects of a cognitive-behavioural internet program on depression, vulnerability to depression and stigma in adolescent males: a school-based controlled trial. *Cogn Behav Ther* 2006; **35**: 43–54.
  - 51 O'Kearney R, Kang K, Christensen H, Griffiths K. A controlled trial of a school-based internet program for reducing depressive symptoms in adolescent girls. *Depress Anxiety* 2009; **26**: 65–72.
  - 52 Ventieri D, Clarke DM, Hay M. The effects of a school-based educational intervention on pre-adolescents' knowledge of and attitudes towards mental illness. *Adv Sch Ment Health Promot* 2011; **4**: 5–17.
  - 53 Gonzales JM, Tinsley HEA, Krauder KR. Effects of psychoeducational interventions on opinions of mental illness, attitudes towards help seeking, and expectations about psychotherapy in college students. *J Coll Stud Dev* 2002; **43**: 51–63.
  - 54 Sharp W, Hargrove DS, Johnson L, Deal WP. Mental health education: an evaluation of a classroom based strategy to modify help seeking for mental health problems. *J Coll Stud Dev* 2006; **47**: 419–38.
  - 55 Faigin DA, Stein CH. Comparing the effects of live and video-taped theatrical performance in decreasing stigmatization of people with serious mental illness. *J Ment Health* 2008; **17**: 594–606.
  - 56 O'Reilly CL, Bell JS, Kelly PJ, Chen TF. Impact of mental health first aid training on pharmacy students' knowledge, attitudes and self-reported behaviours: a controlled trial. *Aust NZ J Psychiatry* 2011; **45**: 549–57.
  - 57 Blair Irvine A, Billow MB, Eberhage MG, Seeley JR, McMahon E, Bourgeois M. Mental illness training for licensed staff in long-term care. *Issues Ment Health Nurs* 2012; **33**: 181–94.
  - 58 Patterson P, Whittington R, Bogg J. Testing the effectiveness of an educational intervention aimed at changing attitudes to self-harm. *J Psychiatr Ment Health Nurs* 2007; **14**: 100–5.
  - 59 Treloar AJ. Effectiveness of education programs in changing clinicians' attitudes toward treating borderline personality disorder. *Psychiatr Serv* 2009; **60**: 1128–31.
  - 60 Jorm AF, Kitchener BA, O'Kearney R, Dear KB. Mental health first aid training of the public in a rural area: a cluster randomized trial. *BMC Psychiatry* 2004; **4**: 33.
  - 61 Fung KMT, Tsang HWH, Cheung WM. Randomized controlled trial of the self-stigma reduction program among individuals with schizophrenia. *Psychiatr Res* 2011; **189**: 208–14.
  - 62 Gumley A, Karatzias A, Power K, Reilly J, McNay L, O'Grady M. Early intervention for relapse in schizophrenia: impact of cognitive behavioural therapy on negative beliefs about psychosis and self-esteem. *Br J Clin Psychol* 2006; **45**: 247–60.
  - 63 Jorm AF, Kitchener BA, Sawyer MG, Scales H, Cvetkovski S. Mental health first aid training for high school teachers: a cluster randomized trial. *BMC Psychiatry* 2010; **10**: 51.

**Table DS1. Medline Search Strategy**

1. Stereotyping/ or Social Stigma/ or Prejudice/ or Attitude/
2. stigma.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]
3. stigma\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]
4. stereotyp\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]
5. Social Perception/
6. Interpersonal Relations/
7. Public Opinion/
8. "Attitude of Health Personnel"/
9. Social Distance/
10. "Rejection (Psychology)"/
11. Human Rights/
12. Civil Rights/
13. Social Justice/
14. public opinion.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]
15. social distance.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]
16. human rights.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]
17. civil rights.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]
18. social justice.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]
19. empower\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]
20. "Discrimination (Psychology)"/
21. discrimination.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]
22. marginali\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword

heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

23. injustice.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

24. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23

25. Mental Health/

26. Mental Disorders/

27. Mentally Ill Persons/

28. Somatoform Disorders/

29. Depression/ or Depression, Postpartum/

30. common mental disorders.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

31. Depressive Disorder/

32. Depressive Disorder, Major/

33. Psychotic Disorders/ or Affective Disorders, Psychotic/

34. Neurotic Disorders/ or Self-Injurious Behavior/ or Anxiety/

35. Anxiety Disorders/

36. Schizophrenia, Paranoid/ or Schizophrenia/

37. Bipolar Disorder/

38. Mood Disorders/

39. Obsessive-Compulsive Disorder/

40. Personality Disorders/

41. Eating Disorders/

42. Anorexia Nervosa/ or Anorexia/

43. Bulimia Nervosa/ or Bulimia/

44. Mental Health Services/

45. mental health.mp.

46. mental illness.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

47. psychotic.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

48. psychosis.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

49. schizo\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

50. bipolar disorder.mp. [mp=title, abstract, original title, name of substance word, subject heading word,

keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

51. manic depression.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

52. OCD.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

53. obsessive compulsive disorder.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

54. anorex\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

55. bulimi\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

56. 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34 or 35 or 36 or 37 or 38 or 39 or 40 or 41 or 42 or 43 or 44 or 45 or 46 or 47 or 48 or 49 or 50 or 51 or 52 or 53 or 54 or 55

57. randomized controlled trial.pt.

58. controlled clinical trial.pt.

59. evaluation studies.pt.

60. Prospective Studies/

61. Follow-Up Studies/

62. Social Marketing/

63. National Health Programs/

64. Government Programs/

65. Program Development/

66. Mass Media/

67. Learning/ or Multimedia/ or Teaching/

68. Internet/

69. Journalism/

70. Video Games/ or Video-Audio Media/ or Video Recording/

71. Software/

72. Pamphlets/

73. Advertising as Topic/

74. Public Relations/

75. Public Relations/

76. Persuasive Communication/

77. Famous Persons/

78. Health Knowledge, Attitudes, Practice/

79. Awareness/

80. Consumer Advocacy/ or Patient Advocacy/

81. Health Promotion/

82. evaluat\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

83. intervention\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

84. Intervention Studies/

85. Randomized Controlled Trials as Topic/

86. randomi\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

87. follow-up.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

88. pre-post.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

89. time series.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

90. time point.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

91. campaign.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

92. message.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

93. advert\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

94. marketing.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

95. public relation\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

96. program\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

97. newspaper\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

98. magazine\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

99. newsletter\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword

heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

100. broadcast.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

101. radio.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

102. television.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

103. cinema.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

104. DVD.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

105. website.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

106. compact disc.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

107. movie.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

108. film.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

109. trailer.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

110. stigma resilience.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

111. stigma change.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

112. (stigma\* adj5 cop\*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

113. (stigma\* adj5 reduc\*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

114. (discrim\* adj5 reduc\*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

115. (stigma\* adj5 challeng\*).mp. [mp=title, abstract, original title, name of substance word, subject heading

word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

116. (discrim\* adj5 challeng\*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

117. (stigma\* adj5 combat\*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

118. (discrim\* adj5 combat\*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

119. (stigma\* adj5 counter\*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

120. (discrim\* adj5 counter\*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

121. contact.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

122. testimon\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

123. mental health literacy.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier]

124. 57 or 58 or 59 or 60 or 61 or 62 or 63 or 64 or 65 or 66 or 67 or 68 or 69 or 70 or 71 or 72 or 73 or 74 or 75 or 76 or 77 or 78 or 79 or 80 or 81 or 82 or 83 or 84 or 85 or 86 or 87 or 88 or 89 or 90 or 91 or 92 or 93 or 94 or 95 or 96 or 97 or 98 or 99 or 100 or 101 or 102 or 103 or 104 or 105 or 106 or 107 or 108 or 109 or 110 or 111 or 112 or 113 or 114 or 115 or 116 or 117 or 118 or 119 or 120 or 121 or 122 or 123

125. 24 and 56 and 124

126. 124 and "Humans".sa\_suba.

127. 125 and 1980:2013.(sa\_year).

128. 126 and 1990:2013.(sa\_year).

129. 127 and 2000:2013.(sa\_year).

## Table DS . References for included studies

- Aho-Mustonen K, Tiihonen J, Repo-Tiihonen E, Ryyanen O-P, Miettinen R, Raty H. Group psychoeducation for long-term offender patients with schizophrenia: An exploratory randomised controlled trial. *Criminal Behaviour and Mental Health*. 2011 Jul;21(3):163-76.
- Altindag A, Yanik M, Ucok A, Alptekin K, Ozkan M. Effects of an antistigma program on medical students' attitudes towards people with schizophrenia. *Psychiatry & Clinical Neurosciences*. 2006;60(3):283-8.
- Anderson K, Austin JC. Effects of a documentary film on public stigma related to mental illness among genetic counselors. *Journal of Genetic Counseling*. 2012 Aug;21(4):573-81.
- Armstrong G, Kermode M, Raja S, Suja S, Chandra P, Jorm AF. A mental health training program for community health workers in India: Impact on knowledge and attitudes. *Int J Ment Health Syst*. 2011 Aug;5:17.
- Bayar MR, Poyraz BC, Aksoy-Poyraz C, Arikan MK. Reducing mental illness stigma in mental health professionals using a web-based approach. *Israel Journal of Psychiatry & Related Sciences*. 2009;46(3):226-30.
- Buizza C, Pioli R, Rossi G, Poli R, Spinogatti F, Agrimi E. Measuring of the effectiveness of an educational intervention in stigma reduction: A pilot study in Brescia and Cremona towns. *Psichiatria e Psicoterapia*. 2007 Sep;26(3):247-58.
- Campbell M, Shryane N, Byrne R, Morrison AP. A mental health promotion approach to reducing discrimination about psychosis in teenagers. *Psychosis: Psychological, Social and Integrative Approaches*. 2011 Feb;3(1):41-51.
- Castro CA, Adler AB, McGurk D, Bliese PD. Mental health training with soldiers four months after returning from Iraq: Randomization by platoon. *Journal of Traumatic Stress*. 2012 Aug;25(4):376-83.
- Clement, S., Van, N. A., Kassam, A., Flach, C., Lazarus, A., De, C. M., Mccrone, P., Norman, I. & Thornicroft, G. 2012. Filmed v. live social contact interventions to reduce stigma: randomised controlled trial. *Br.J.Psychiatry*, 201, 57-64.
- Conrad I, Dietrich S, Heider D, Blume A, Angermeyer MC, Riedel-Heller S. "Crazy? So what!": A school programme to promote mental health and reduce stigma-Results of a pilot study. *Health Education*. 2009;109(4):314-28.
- Corker E. HS, Henderson C., Weeks C., Pinfold V., Rose D., Williams P., Flach C., Gill V. & Thornicroft. Experiences of discrimination among people using mental health services in England 2008-11. *British Journal of Psychiatry*. 2013 2013;202(55):s58-s63.
- Crisp A, Gelder M, Goddard E, Meltzer H. Stigmatization of people with mental illnesses: a follow-up study within the Changing Minds campaign of the Royal College of Psychiatrists. *World Psychiatry*. 2005 Jun;4(2):106-13.
- Dietrich S, Mergl R, Freudenberg P, Althaus D, Hegerl U. Impact of a campaign on the public's attitudes towards depression. *Health Education Research*. 2010;25(1):135-50.
- Domino G. Impact of the film 'One flew over the cuckoo's nest' on attitudes towards mental illness. *Psychological Reports*. 1983 1983/08/01;53(1):179-82.
- Economou M, Louki E, Peppou L, Gramandani C, Yotis L, Stefanis C. Fighting psychiatric stigma in the classroom: The impact of an educational intervention on secondary school students' attitudes to schizophrenia. *International Journal of Social Psychiatry*. 2012 Sep;58(5):544-51.
- Essler V, Arthur A, Stickley T. Using a school-based intervention to challenge stigmatizing attitudes and promote mental health in teenagers. *Journal of Mental Health*. 2006;15(2):243-50.

Esters IG, Cooker PG, Ittenbach RF. Effects of a unit of instruction in mental health on rural adolescents' conceptions of mental illness and attitudes about seeking help. *Adolescence*. 1998;33(130):469-76.

Evans-Lacko S, Henderson C, Thornicroft G, McCrone P. Economic evaluation of the anti-stigma social marketing campaign in England 2009-2011. *Br J Psychiatry Suppl*. 2013 Apr;55:s95-101.

Evans-Lacko S, Malcolm E, West K, Rose D, London J, Rusch N, et al. Influence of Time to Change's social marketing interventions on stigma in England 2009-2011. *Br J Psychiatry Suppl*. 2013 Apr;55:s77-88.

Faigin DA, Stein CH. Comparing the effects of live and video-taped theatrical performance in decreasing stigmatization of people with serious mental illness. *Journal of Mental Health*. 2008;17(6):594-606.

Friedrich B, Evans-Lacko S, London J, Rhydderch D, Henderson C, Thornicroft G. Anti-stigma training for medical students: the Education Not Discrimination project. *Br J Psychiatry Suppl*. 2013 Apr;55:s89-94.

Fung KMT, Tsang HWH, Cheung W-m. Randomized controlled trial of the self-stigma reduction program among individuals with schizophrenia. *Psychiatry Research*. 2011 Sep 30;189(2):208-14.

Gaebel W, Zaska H, Baumann AE, Klosterkötter J, Maier W, Decker P, et al. Evaluation of the German WPA "Program against stigma and discrimination because of schizophrenia--Open the Doors": Results from representative telephone surveys before and after three years of antistigma interventions. *Schizophrenia Research*. 2008 Jan;98(1-3):184-93.

Garcia Silberman S. Mental health and the movies: An experimental study about attitude change. *Revista Mexicana de Psicología*. 1993 Jan-Jun;10(1):37-48.

Ghorbani NR, Heidari RN. Effects of information and communication technology on youth's health knowledge. *Asia-Pacific Journal of Public Health*. 2011 May;23(3):363-8.

Gonzales JM, H.E.A. T, K.R. K. Effects of psychoeducational interventions on opinions of mental illness, attitudes towards help seeking, and expectations about psychotherapy in college students. *Journal of College Student Development*. 2002;43:51-63.

Gould M, Greenberg N, Hetherington J. Stigma and the military: evaluation of a PTSD psychoeducational program. *JTrauma Stress*. [10.1002/jts.20233 doi]. 2007;20(4):505-15.

Graham AL, Julian J, Meadows G. Improving responses to depression and related disorders: Evaluation of a innovative, general, mental health care workers training program. *Int J Ment Health Syst*. 2010 Sep;4:25.

Grausgruber A, Schony W, Grausgruber-Berner R, Koren G, Apor BF, Wancata J, et al. ["Schizophrenia has many faces" - evaluation of the Austrian Anti-Stigma-Campaign 2000-2002]. *Psychiatrische Praxis*. [English Abstract]. 2009 Oct;36(7):327-33.

Griffiths, K. M., Carron-Arthur, B., Parsons, A. & Reid, R. 2014. Effectiveness of programs for reducing the stigma associated with mental disorders. A meta-analysis of randomized controlled trials. *World Psychiatry* : , 13, 161-75.

Gulliver A, Griffiths KM, Christensen H, Mackinnon A, Calear AL, Parsons A, et al. Internet-based interventions to promote mental health help-seeking in elite athletes: An exploratory randomized controlled trial. *Journal of Medical Internet Research*. 2012 May-Jun;14(3):120-37.

Gumley A, Karatzias A, Power K, Reilly J, McNay L, O'Grady M. Early intervention for relapse in schizophrenia: Impact of cognitive behavioural therapy on negative beliefs about psychosis and self-esteem. *British Journal of Clinical Psychology*. 2006 Jun;45(2):247-60.

Gutierrez-Maldonado J, Caqueo-Urizar A, Ferrer-Garcia M. Effects of a psychoeducational intervention program on the attitudes and health perceptions of relatives of patients with schizophrenia. *Social Psychiatry and Psychiatric Epidemiology*. 2009 May;44(5):343-8.

Hao Z-h, Liang B-y. Effect of group guidance on intention of seeking professional psychological help among university students. *Chinese Journal of Clinical Psychology*. 2011 Oct;19(5):699-702.

Henderson C, Corker E, Lewis-Holmes E, Hamilton S, Flach C, Rose D, et al. England's time to change antistigma campaign: one-year outcomes of service user-rated experiences of discrimination. *Psychiatric Services*. . 2012;63(5):451-7.

Henderson C, Williams P, Little K, Thornicroft G. Mental health problems in the workplace: changes in employers' knowledge, attitudes and practices in England 2006-2010. *Br J Psychiatry Suppl*. 2013 Apr;55:s70-6.

Irvine BA, Billow MB, Eberhage MG, Seeley JR, McMahon E, Bourgeois M. Mental illness training for licensed staff in long-term care. *Issues in Mental Health Nursing*. 2012 Mar;33(3):181-94.

Jorm AF, Christensen H, Griffiths KM. Changes in depression awareness and attitudes in Australia: The impact of beyondblue: The national depression initiative. *Australian and New Zealand Journal of Psychiatry*. 2006 Jan;40(1):42-6.

Jorm AF, Christensen H, Griffiths KM. The impact of beyondblue: the national depression initiative on the Australian public's recognition of depression and beliefs about treatments. *Australian and New Zealand Journal of Psychiatry*. 2005;39(4):248-54.

Jorm AF, Kitchener BA, Fischer J-A, Cvetkovski S. Mental health first aid training by e-learning: A randomized controlled trial. *Australian and New Zealand Journal of Psychiatry*. 2010 Dec;44(12):1072-81.

Jorm AF, Kitchener BA, O'Kearney R, Dear KB. Mental health first aid training of the public in a rural area: A cluster randomized trial. *BMC Psychiatry*. 2004 Oct;4:33.

Jorm AF, Kitchener BA, Sawyer MG, Scales H, Cvetkovst S. Mental health first aid training for high school teachers: a cluster randomised trial. *BMC Psychiatry* 2010 10:51

Kelly CM, Mithen JM, Fischer JA, Kitchener BA, Jorm AF, Lowe A, et al. Youth mental health first aid: a description of the program and an initial evaluation. *Int J Ment Health Syst*. 2011 Jan;5.

Kerby J, Calton T, Dimambro B, Flood C, Glazebrrok C. Anti-stigma films and medical students' attitudes towards mental illness and psychiatry: randomised controlled trial. *Psychiatric Bulletin*. 2008;32:345-9.

Kitchener BA, Jorm AF. Mental health first aid training for the public: evaluation of effects on knowledge, attitudes and helping behavior. *BMC Psychiatry*. 2002;2(1):10.

Kitchener BA, Jorm AF. Mental health first aid training in a workplace setting: A randomized controlled trial. *BMC Psychiatry*. 2004 Aug;4:23.

Knight MT, Wykes T, Hayward P. Group Treatment of Perceived Stigma and Self-Esteem in Schizophrenia: A Waiting List Trial of Efficacy. *Behavioural and Cognitive Psychotherapy*. 2006 Jul;34(3):305-18.

Krawitz R. Borderline personality disorder: attitudinal change following training. *Australian & New Zealand Journal of Psychiatry*. 2004;38(7):554-9.

Link BG, Struening EL, Neese-Todd S, Asmussen S, Phelan JC. On Describing and Seeking to Change the Experience of Stigma. [References]. *Psychiatric Rehabilitation Skills*. 2002;6(2):231.

Luty J, Rao H, Arokiadass SMR, Easow JM, Sarkhel A. The repentant sinner: Methods to reduce stigmatised attitudes towards mental illness.

Psychiatric Bulletin. 2008 Sep;32(9):327-32.

Masuda A, Hayes SC, Lillis J, Bunting K, Herbst SA, Fletcher LB. The relation between psychological flexibility and mental health stigma in Acceptance and Commitment Therapy: A preliminary process investigation. *Behavior and Social Issues*. 2009 Spr-Sum;18(1):1-16.

Mehta N, Kassam A, Leese M, Butler G, Thornicroft G. Public attitudes towards people with mental illness in England and Scotland, 1994-2003. *Br J Psychiatry*. [Research Support, Non-U.S. Gov't]. 2009 Mar;194(3):278-84.

Naylor PB, Cowie HA, Walters SJ, Talamelli L, Dawkins J. Impact of a mental health teaching programme on adolescents. *British Journal of Psychiatry*. 2009 Apr;194(4):365-70.

Ng P, Chan KF. Attitudes towards people with mental illness. Effects of a training program for secondary school students. *International journal of adolescent medicine and health*. 2002;14(3):215-24.

O'Reilly CL, Bell JS, Kelly PJ, Chen TF. Impact of mental health first aid training on pharmacy students' knowledge, attitudes and self-reported behaviours: a controlled trial. *Australia and New Zealand Journal of Psychiatry* 2011 45: 549-557

O'Kearney R, Gibson M, Christensen H, Griffiths KM. Effects of a cognitive-behavioural internet program on depression, vulnerability to depression and stigma in adolescent males: A school-based controlled trial. *Cognitive Behaviour Therapy*. 2006;35(1):43-54.

O'Kearney R, Kang K, Christensen H, Griffiths K. A controlled trial of a school-based Internet program for reducing depressive symptoms in adolescent girls. *Depression and Anxiety*. 2009;26(1):65-72.

O'Reilly CL, Bell JS, Chen TF. Consumer-led Mental Health Education for Pharmacy Students. *American Journal of Pharmaceutical Education*. 2010;74(9).

Patterson P, Whittington R, Bogg J. Testing the effectiveness of an educational intervention aimed at changing attitudes to self-harm. *Journal of Psychiatric & Mental Health Nursing*. 2007;14(1):100-5.

Paykel ES, Hart D, Priest RG. Changes in public attitudes to depression during the Defeat Depression Campaign. *Br J Psychiatry*. 1998;173:519-22.

Pejovic-Milovancevic M, Lecic-Tosevski D, Tenjovic L, Popovic-Deusic S, Draganic-Gajic S. Changing attitudes of high school students towards peers with mental health problems. *PsychiatrDanub*. 2009;21(2):213-9.

Pierce D, Liaw S-T, Dobell J, Anderson R. Australian rural football club leaders as mental health advocates: An investigation of the impact of the Coach the Coach project. *Int J Ment Health Syst*. 2010 May;4:10.

Pinfold V, Huxley P, Thornicroft G, Farmer P, Toulmin H, Graham T. Reducing psychiatric stigma and discrimination--evaluating an educational intervention with the police force in England. *Social Psychiatry and Psychiatric Epidemiology*. 2003;38(6):337-44.

Pinfold V, Toulmin H, Thornicroft G, Huxley P, Farmer P, Graham T. Reducing psychiatric stigma and discrimination: evaluation of educational interventions in UK secondary schools. *Br J Psychiatry*. 2003;182:342-6.

Pinto-Foltz MD, Logsdon MC, Myers JA. Feasibility, acceptability, and initial efficacy of a knowledge-contact program to reduce mental illness stigma and improve mental health literacy in adolescents. *Social Science and Medicine*. [S0277-9536(11)00242-5 pii ;10.1016/j.socscimed.2011.04.006 doi]. 2011;72(12):2011-9.

Premalatha Chinnayya H, Chandrashekar CR, Moily S, Puttamma, Raghuram A, Subramanya KR, et al. Training Primary Care Health Workers in Mental Health Evaluation of Attitudes towards Mental Illness before and after Training. *International Journal of Social Psychiatry*. 1990 December 1, 1990;36(4):300-7.

Rahman A, Mubbashar MH, Gater R, Goldberg D. Randomised trial of impact of school mental-health programme in rural Rawalpindi, Pakistan. *Lancet*. 1998;352(9133):1022-5.

Reavley NJ, Jorm AF. Stigmatising attitudes towards people with mental disorders: Changes in Australia over 8 years. *Psychiatry Research*. 2012 May;197(3):302-6.

Rong Y, Glozier N, Luscombe GM, Davenport TA, Huang Y, Hickie IB. Improving knowledge and attitudes towards depression: A controlled trial among Chinese medical students. *BMC Psychiatry*. 2011 Mar;11:36.

Schneider J, Beeley C, Repper J. Campaign appears to influence subjective experience of stigma. *Journal of Mental Health*. 2011 Feb;20(1):89-97.

Schulze B, Richter-Werling M, Matschinger H, Angermeyer MC. Crazy? So what! Effects of a school project on students' attitudes towards people with schizophrenia. *Acta Psychiatr Scand*. 2003;107(2):142-50.

Seal KH, Abadjian L, McCamish N, Shi Y, Tarasovsky G, Weingardt K. A randomized controlled trial of telephone motivational interviewing to enhance mental health treatment engagement in Iraq and Afghanistan veterans. *General Hospital Psychiatry*. 2012 Sep;34(5):450-9.

Shandley K, Austin D, Klein B, Kyrios M. An evaluation of 'Reach Out Central': an online gaming program for supporting the mental health of young people. *Health Education Research*. 2010;25(4):563-74.

Sharp W, Hargrove DS, Johnson L, Deal WP. Mental Health Education: An Evaluation of a Classroom Based Strategy to Modify Help Seeking for Mental Health Problems. *Journal of College Student Development*. 2006 Jul-Aug;47(4):419-38.

Shi Zhongying, Liu Shaoyan, Nie Leiyan The impact of group psychological interventions on the stigma perceived of patients with depression. *Journal of Nurses Training* 2012, 27:1213-1216.

Sogaard AJ, Fonnebo V. The Norwegian Mental Health Campaign in 1992. Part II: changes in knowledge and attitudes. *Health Education Research*. 1995;10(3):267-78.

Treloar AJ. Effectiveness of education programs in changing clinicians' attitudes toward treating borderline personality disorder. *Psychiatric Services*. 2009;60(8):1128-31.

Ucok A, Soygur H, Atakli C, Kuscu K, Sartorius N, Duman ZC, et al. The impact of antistigma education on the attitudes of general practitioners regarding schizophrenia. *Psychiatry and Clinical Neurosciences*. 2006 Aug;60(4):439-43.

Ventieri D, Clarke DM, Hay M. The effects of a school-based educational intervention on pre-adolescents' knowledge of and attitudes towards mental illness. *Advances in School Mental Health Promotion*. 2011 Jul;4(3):5-17.

Worakul P, Thavichachart N, Lueboonthavatchai P. Effects of psycho-educational program on knowledge and attitude upon schizophrenia of schizophrenic patients' caregivers. *J Med Assoc Thai*. 2007 2007;90(6):1199-204.

Wright A, McGorry PD, Harris MG, Jorm AF, Pennell K. Development and evaluation of a youth mental health community awareness campaign - The Compass Strategy. *BMC Public Health*. 2006;6(215).

Yamaguchi S, Mino Y. [Effects of an educational intervention to reduce prejudice toward the mentally disabled among high school students]. [Nihon koshu eisei zasshi] *Japanese journal of public health*. 2007;54(12):839-46.

Yanos PT, Roe D, West ML, Smith SM, Lysaker PH. Group-based treatment for internalized stigma among persons with severe mental illness: Findings from a randomized controlled trial. *Psychological Services*. 2012 Aug;9(3):248-58.



**Table DS3. Medium/long-term effectiveness of interventions to reduce stigma: study characteristics, statistical significance and direction of findings (RCTs, and con**

Study	Intervention Scale	Country	HIC or LAMIC <sup>a</sup>	Design <sup>b</sup>	Participant type	Intervention Group (n)	Control Group (n)	Intervention Description	Disorder focused on	Control description
<b>Targeted at the Armed Forces</b>										
Seal 2012	Target population	USA	HIC	RCT	Middle east war veterans screened positive for mental health problems	34	39	Four unscripted 20-30 min telephone sessions of motivational interviewing.	Mental illness	Consisted in providing participants with mental health assessment results and advising about the logistics of treatment referrals, e.g. clinic hours, parking, transportation etc.
Castro 2012	Target population	USA	HIC	RCT	US soldiers	804	841	Mental health training program developed for military personnel across the deployment cycle.	Stress and PTSD	No intervention
Gould 2007	Target population	UK	HIC	Controlled <sub>1</sub>	Active service members of the UK Armed Forces	62	62	Training programme which seeks to modify attitudes about PTSD, provide support and education, and identify and refer at risk personnel.	PTSD	No intervention
<b>Targeted at School Students</b>										
Campbell 2011	Target population	England	HIC	RCT	Students (14-15 years)	43	49	Mental health workshop including an interactive educational presentation and a personal testimony of psychotic experiences by a service user researcher.	Mental illness	No intervention
Pinto-Foltz 2011	Target population	USA	HIC	RCT	Female students (13-17 years)	95	61	Narrative story-telling, discussion and a video presentation administered by two trained service users who were in sustained recovery from mental illness.	Mental illness	No intervention
Conrad 2009	Target population	Germany	HIC	Controlled <sub>1</sub>	Students (13-18 years)	120	90	Introduction to mental health, discussion about students' perceptions of life and a discussion with other young people with lived experience mental illness.	Mental illness	No intervention
Economou 2012	Target population	Greece	HIC	Controlled <sub>2</sub>	Students (13-15 years)	308	308	A two hour session including a discussion of 'myths' regarding schizophrenia, patient narratives with described experiences of stigma and a role-play session.	Schizophrenia	A talk about nutrition and healthy living
Esters 1998	Target population	USA	HIC	Controlled <sub>1</sub>	Students enrolled in health classes (13 to 17 years)	20	20	Three day programme including a video and information about sources of help for mental health problems and the reality of associated stigma.	Mental illness	Regularly scheduled classes unrelated to mental health
Naylor 2009	Target population	England	HIC	Controlled <sub>1</sub>	Students (14-15 years)	174	242	Six lessons on mental health issues delivered over a six week period.	Mental illness	Different school in which intervention did not occur (inactive)
Ng 2002	Target population	China (Hong Kong)	HIC	Controlled <sub>1</sub>	Students (mean age: 14-15 years)	79	90	School based mental health club including a weekly 1h mental health education program, a mental health promotion day, talks and exhibitions, and direct contact with people with mental health problems.	Mental illness	No intervention
O'Kearney 2006	Target population	Australia	HIC	Controlled <sub>1</sub>	Male Students (15-16 years)	35	24	An interactive Internet programme aiming to help people identify, overcome and cope with depression. Skills and knowledge based with information, demonstrations, questionnaires and practice exercises.	Depression	Usual personal development programme scheduled by the school consisting in private study, ad hoc discussion and physical activities with no specific discussion of depression.
O'Kearney 2009	Target population	Australia	HIC	Controlled <sub>1</sub>	Female Students (15-16 years)	67	90	An interactive Internet programme aiming to help people identify, overcome and cope with depression. Skills and knowledge based with information, demonstrations, questionnaires and practice exercises.	Depression	School's normal personal development activities about nutrition.

Schulze 2003	Target population	Germany	HIC	Controlled <sub>1</sub>	Students (14-18 years)	90	60	A week dedicated to an anti-stigma initiative including discussions of wellbeing, mental illness, responses to low mood and causes of illness, a personal testimony from a young person with schizophrenia and further discussions about schizophrenia and stigma.	Schizophrenia	1 week of a different unrelated project
Ventieri 2011	Target population	Australia	HIC	Controlled <sub>1</sub>	Students (9-12 years)	69	126	Educational intervention including a general introduction to mental illness, causes and treatments of mental illness and stigma.	Mental illness	Normal classes covering material unrelated to mental illness
Yamaguchi 2007 <sup>2</sup>	Target population	Japan	HIC	Controlled <sub>1</sub>	Students (16-18 years)	99	81	50-min mental health lecture and 50-min patient's talk at class homeroom	Mental illness	No intervention
Buizza 2007 (Brescia)	Target population	Italy	HIC	Uncontrolled	Students (16-19 years)	327	n/a	A 2 hour session including a stigma video, an educational talk about mental health illnesses and stigma, a live testimony by a mental health service user (or family member) and a final discussion.	Mental illness	n/a
Buizza 2007 (Cremona)	Target population	Italy	HIC	Uncontrolled	Students (16-19 years)	380	n/a	A 2 hour session including a stigma video, an educational talk about mental health illnesses and stigma, a live testimony by a mental health service user (or family member) and a final discussion.	Mental illness	n/a
Essler 2006	Target population	England	HIC	Uncontrolled	Students (13-14 years)	104	n/a	Intervention delivered by a theatre company aimed at increasing knowledge about mental health issues, promoting positive attitudes towards people with mental health problems, and increasing awareness of the pupils' own attitudes.	Mental illness	n/a
Pejovic-Milovancevic 2009	Target population	Serbia	LAMIC	Uncontrolled	Students (15 years)	63	n/a	6 week anti-stigma programme including educational sessions covering mental health problems, discrimination, common myths and a weekly interactive workshop where targeted activities were carried out.	Mental illness	n/a
Pinfold 2003	Target population	UK	HIC	Uncontrolled	Students (14-15 years)	472	n/a	Two hour educational intervention including mental health awareness workshops, a video about people living with schizophrenia and discussions about labels and stereotypes or a session co-facilitated by a person who had experiences of living with mental health problems.	Mental illness	n/a
<b>Targeted at University Students</b>										
Gonzales 2002	Target population	USA	HIC	RCT	Psychology University Students	167 for both interventions and controls	167 for both interventions and controls	Educational intervention in which participants read information about mental illness and stigma.	Mental illness	No intervention
Kerby 2008	Target population	UK	HIC	RCT	Medical Students	23	23	2 films produced by mental health service users combining both education and stereotype disconfirmation elements.	Mental illness	25 minute documentary unrelated to mental illness or psychiatry, matched for visual format.
Sharp 2006	Target population	USA	HIC	RCT	University Students	62	61	A 40 minute lecture including information about psychological disorders, the therapeutic process, common misconceptions about psychotherapy and information about local mental health services.	Mental illness	The control group met for 40 minutes and watched an astronomy science video.
Domino 1983	Target population	USA	HIC	Controlled <sub>1</sub>	University Students	37	19	Watching the film 'One Flew Over The Cuckoo's Nest' in a public cinema.	Mental illness	Not having watched the film

Faigin 2008	Target population	USA	HIC	Controlled <sub>1</sub>	Psychology University Students	81	123	A play conducted by actors with history of severe mental illness addressing issues such as group-home living, hospitalization, and the impact of social stigmatization. Included a question and answer session.	Mental illness	No intervention
Faigin 2008 (2nd arm)	Target population	USA	HIC	Controlled <sub>1</sub>	Psychology University Students	99	123	Video group - viewed a recorded version of the play described above.	Mental illness	No intervention
Friedrich 2013	Target population	UK	HIC	Controlled <sub>1</sub>	Medical Students	1066	386	Short lecture with key facts about stigma and discrimination, testimonies about the experiences of mental health problems and stigma from people with direct experience and role-plays in small groups.	Mental illness	No intervention
O'Reilly 2011	Target population	Australia	HIC	Controlled <sub>1</sub>	Pharmacy Students	60	212	Mental Health First Aid training course consisting in mental health lectures, supervised weekly placements in the community pharmacy and a tutorial on a mental health topic.	Mental illness	Group who did not attend the course (because they did not wish to and because there was not enough space available on the course)
Altindag 2006	Target population	Turkey	LAMIC	Controlled <sub>2</sub>	Medical Students	25	35	1 day program with an educational lecture about stigma and schizophrenia, direct contact with a young person with schizophrenia and indirect contact, viewing of the film 'A Beautiful Mind'.	Schizophrenia	1 day program with a 2-h lecture about water metabolism and a viewing of a documentary on the migratory patterns of birds.
Masuda 2009	Target population	USA	HIC	Uncontrolled	Psychology University Students	27	n/a	Acceptance and Commitment Therapy delivered in a 150-minute workshop format. The major goal of the intervention was to humanise the stigmatised individuals.	Mental illness	n/a
O'Reilly 2010	Target population	Australia	HIC	Uncontrolled	Pharmacy Students	258	n/a	Mental health lectures, placements in community pharmacy settings, and a tutorial with trained mental health consumer educators. Students asked the educators about their experience of medication history and counselling.	Mental illness	n/a
Rong 2011	Target population	China	LAMIC	Uncontrolled	Medical Students	103	n/a	The intervention consisted in a 1.5 hour lecture which covered all the basic medical aspects about depression required by the teaching guidelines	Depression	n/a
Rong 2011 (2nd arm)	Target population	China	LAMIC	Uncontrolled	Medical Students	102	n/a	The intervention consisted in a 1.5 hour lecture about depression and 2 weeks of additional activities carried out by students about the experience of mental illness.	Depression	n/a
<b>Targeted at Healthcare Professionals</b>										
Irvine 2012	Target population	USA	HIC	RCT	Licensed care staff in long-term care facilities	84	88	Two Internet courses with video-based behavioural skills and knowledge training.	88	No intervention
Patterson 2007	Target population	UK	HIC	Controlled <sub>2</sub>	Healthcare professionals	69	22	Educational intervention over 12 days which aimed to improve knowledge of causes of self-harm and reflect on feelings and responses to self-harming behaviour.	Self-harm and suicide	Course unconnected to self harm, on research method
Treloar 2009	Target population	Australia and New Zealand	HIC	Controlled <sub>2</sub>	Health practitioners with experience of patients with BPD	49	41	Educational programme which covered research and attitudes about borderline personality disorder and discussions of BPD using case studies.	Borderline personality disorder	No intervention
Treloar 2009	Target population	Australia and New Zealand	HIC	Controlled <sub>2</sub>	Health practitioners with experience of patients with BPD	49	41	A cognitive-behavioural education programme using case studies and a discussion of clinical guidelines for	Borderline personality disorder	No intervention

(2nd arm)	Target population	Zealand	HIC	Controlled <sup>2</sup>	Experience of patients with BPD	n <sub>1</sub>	n <sub>2</sub>	Case studies and a discussion of 01 critical guidelines for self-harm.	Borderline personality disorder	No intervention
Armstrong 2011	Target population	India	LAMIC	Uncontrolled	Community health workers	70	n/a	Mental health training program aiming to increase recognition of mental disorders, enhance appropriate response and referral, support people with mental disorders and their families, and improve mental health promotion in communities.	Mental illness	n/a
Graham 2010	Target population	Australia	HIC	Uncontrolled	Workers from diverse sectors of the mental health workforce	1126	n/a	Mental health aptitudes into practice training package including an introduction to mental health and mental illness, depression and anxiety.	Mental illness	n/a
Krawitz 2004	Target population	Australia	HIC	Uncontrolled	Staff working with BPD patients from public mental health and substance abuse services	418	n/a	2 day educational workshop about BPD including educational media, interactional learning and structured discussion.	Borderline personality disorder	n/a
Ücock 2006	Target population	Turkey	LAMIC	Uncontrolled	General Practitioners	106	n/a	Training session including information about the course of schizophrenia, treatment, the impact of stigma and a description of GP roles followed by an discussion session.	Schizophrenia	n/a
<b>Targeted at the General Public</b>										
Jorm 2010	Large Scale	Australia	HIC	RCT*	General population	90	84	Mental Health First Aid e-learning CD which teaches members of the public to recognize the early signs of mental illness and to provide initial help to a person developing a mental disorder or in a mental health crisis situation delivered with the aid of video clips and interactive case studies.	Mental illness	Waiting list control
Jorm 2010 (2nd arm)	Large Scale	Australia	HIC	RCT*	General population	88	84	Mental Health First Aid manual, containing the same elements as above but in written format only.	Mental illness	Waiting list control
Jorm 2004	Large Scale	Australia	HIC	RCT*	Residents in regional catchment areas	416	337	9 hour Mental Health First Aid course which teaches how to help people in the crisis situations of being suicidal, having a panic attack, being exposed to a traumatic event, or in an acute psychotic state.	Mental illness	Waiting list control
Luty 2008	Large Scale	UK	HIC	RCT	General population	148	155	Vignettes containing a photograph of a professional-looking male model with an 'upbeat' description of a hypothetical patient in remission.	Schizophrenia	Case description of a patient
Kelly 2011	Small Scale	Australia	HIC	Uncontrolled	General public	246	n/a	Youth Mental Health First Aid course which teaches adults how to support adolescents who might be developing a mental health problem/in a mental health crisis and to assist them to receive professional help.	Mental illness in adolescents	n/a
Kitchener 2002	Small Scale	Australia	HIC	Uncontrolled	General public	190	n/a	Mental Health First Aid training (9 hours) which covers how to help people in mental health crises and/or in the early stages of mental health problems and educational information about mental illnesses.	Mental illness	n/a
Shandley 2010	Target population	Australia	HIC	Uncontrolled	General population, females aged 18-25 years	266	n/a	A web-based interactive educational game designed to support the mental health of young people and assisting them in identifying and developing coping skills dealing with precursors to mental health problems.	Mental illness	n/a
Shandley 2010 (2nd arm)	Target population	Australia	HIC	Uncontrolled	General population, males aged 18-25 years	266	n/a	A web-based interactive educational game designed to support the mental health of young people and assisting them in identifying and developing coping skills dealing with precursors to mental health problems.	Mental illness	n/a
Sogaard 1992	Large Scale	Norway	HIC	Uncontrolled	General population	T1 survey: 1191	T2 survey: 574	A 6h TV show consisting in information on mental health problems and prevention interspersed with	Mental illness	n/a

								entertainment and reporting fundraising results.		
<b>Targeted at people with mental health problems</b>										
Fung 2011	Target population	China (Hong Kong)	HIC	RCT*	People with schizophrenia receiving psychiatric services in the community	34	32	Self-stigma reduction programme containing 16 sessions covering psychoeducation, self-esteem, readiness for change, social skills and a goal attainment.	Schizophrenia	Newspaper reading group
Aho-Mustonen 2011	Target population	Finland	HIC	RCT	Long-stay forensic patients of a maximum security forensic psychiatric hospital	19	20	A group psychoeducation programme consisting of eight group sessions which were conducted once a week.	Schizophrenia	Treatment as usual
Gumley 2006	Target population	Wales and Ireland	HIC	RCT	People diagnosed with schizophrenia (or related disorder) prone to relapse	72	72	CBT consisting of a five-session engagement phase delivered and an intensive targeted phase.	Schizophrenia	Treatment as usual (ongoing medication, regular psychiatric review and regular follow-up from key worker)
Link 2002	Target population	USA	HIC	RCT	People with a serious mental illness	88 including controls	88 including controls	16 sessions consisting in specific approaches to developing an awareness of stigma, internalisation of stigma and its consequences, identification of stigma in social interactions and choosing ways to cope with it.	Mental illness	Waiting list control
Yanos 2012	Target population	USA	HIC	RCT	People with a serious mental illness who met criteria for elevated internalized stigma	15	18	Narrative enhancement/cognitive therapy including psychoeducation, cognitive restructuring and narrative enhancement which addresses themes of hope which contrast with stigmatising views.	Mental illness	Treatment as usual (did not include any services which focused on internalised stigma)
Knight 2006	Target population	England	HIC	Waiting list control	People with schizophrenia spectrum disorders	21	n/a	Two CBT interventions conducted over 7 weekly sessions focused on educating and emphasizing the similarity of experience of group members, self-esteem, and highlighting maladaptive coping strategies. Included psychoeducation about stigma and mental illness.	Schizophrenia	6 week waiting control in which participants were assessed prior to the intervention.
Schneider 2011	Small Scale	England	HIC	Uncontrolled	Service users of an English mental health trust	243	n/a	Anti-stigma campaign in an NHS trust for one month which aimed to improve public attitudes, remove barriers to service users' working, promote recovery and social inclusion and reduce the incidence of service users' experiences of discrimination and self-stigma.	Mental illness	n/a
<b>Other</b>										
Gulliver 2012	Target population	Australia	HIC	RCT*	Elite Athletes	30	29	Mental Health literacy and destigmatization intervention including online educational material about common mental health disorders, stigma and help-seeking sources.	Mental illness	No intervention
Gulliver 2012 (2nd arm)	Target population	Australia	HIC	RCT*	Elite Athletes	30	29	Comprised interactive quizzes providing tailored feedback to the participant about his or her level of depression and anxiety allowing the participant to assess their need to seek help.	Mental illness	No intervention
Gulliver 2012 (3rd arm)	Target population	Australia	HIC	RCT*	Elite Athletes	31	29	Help-seeking intervention including an online list of help seeking sources and a congratulatory page.	Mental illness	No intervention

Kitchener 2004	Target population	Australia	HIC	RCT*	Government Employees	146	155	9 hour Mental Health First Aid course which teaches how to help people in mental health crises and/or in the early stages of mental health problem. Includes mental health education about depressive, anxiety and psychotic disorders.	Mental illness	Waiting list control
Jorm 2010	Target population	Australia	HIC	RCT	School teachers of pupils aged 12-15 years	221	106	Youth Mental Health First Aid course completed over 2 days including departmental policy on mental health issues, common mental disorders in adolescents and how to apply the mental health action plan to help a students with such problem.	Mental illness	Waiting list control
Anderson 2012	Target population	USA and Canada	HIC	Uncontrolled	Certified genetic counsellors and genetic counselling students	87	n/a	A 45 minute documentary film "Cracking Up" that follows a group of individuals with mental illness as they participate in a programme that teaches individuals with mental illness how to perform stand-up comedy.	Mental illness	n/a
Pierce 2010	Target population	Australia	HIC	Uncontrolled	Football Coaches	36	n/a	Mental health first training covering depressive, anxiety and psychotic disorders, with education of overlapping substance abuse for 12 hours over 3 weeks.	Mental illness	n/a
Pinfold 2003	Target population	England	HIC	Uncontrolled	Police Officers	163	n/a	An educational intervention delivered by service users, carers and people working in the field of mental health. Included a hearing voices simulation exercise, personal testimonies of psychotic experiences and recovery, and 'How can the police support people with mental health problems?'.	Mental illness	n/a

a High income country (HIC) or low to middle income country (LAMIC). Based on World Bank classifications as of July 1, 2012

b In studies which measured outcomes at more than 2 time points, only the first and last assessments were included in the table. In RCTs, only time\*intervention interactions were reported. Where this was carried across multiple time points, only the final time point results were reported. When studies included between and within-group analyses, only the between-group analyses were reported.

c + indicates p< 0.05 in a direction of benefit (reduction in stigma); 0 indicates p>0.05; - indicates p<0.05 in direction of harm (increase in stigma). Each symbol represents one outcome (scales, or item)











	Item				Attitudes Towards Seeking Professional Psychological Help Scale - Short Form (ATSPPH-SF) (Fischer & Farina, 1995) Intentions Scale of the General Help-seeking Questionnaire (GHSQ) (Deane et al, 2001) Actual Help-seeking Questionnaire (AHSQ) (Wilson et al, 2002)						
5 months	Scale	00	+		Recognition of Mental Illness using vignettes (Kitchener et al., 2002) Beliefs about Treatment for Depression and Schizophrenia (vignettes) (Kitchener et al., 2002) Social Distance Scale (Link et al., 1999)						
	Item	00		0	Contact with someone with a mental health problem (Kitchener et al., 2002)						
6 months	Scale	++			Mental Health Literacy (Jorm et al, 1997) Perceived Stigma Scale (Griffiths, 2008) Personal Stigma Scale (Griffiths et al, 2008)						
	Item	0000000+	000000+	0++++	School Policies on Student Mental Health (developed by authors)						
1 month	Scale	0	0		Stereotype Endorsement scale (Nordt et al., 2006) Social Distance Scale (Penn et al., 1994)						
	Item										
6 months	Scale				Recognition of Depression and Schizophrenia (vignettes) (Kitchener & Jorm, 2004) Responses to Depression (Kitchener & Jorm, 2004) Depression Management (Kitchener & Jorm, 2004)						
	Item	00000000000000 00+++++									
4 weeks	Scale	0	0+++		Knowledge of mental illness (Pinfold et al., 2003a) Officers endorsements of 5 key workshop messages (Pinfold et al., 2003a)						
	Item	+++++			Community Attitudes towards Mental Illness (Taylor and Dear, 1981) and WPA Alberta Pilot site questionnaire tool kit (WPA, 2000) used to create an Attitude measure (3 scales) and a Social Distance measure (4 items)						

analyses, only the between group data was extracted. In studies with a control group where data was analysed within-group, results were only reported as a benefit or harm when no changes were detected in the control group. In pre-post studies with a control group, we only reported findings at time points where control data was present.

When a study included unadjusted and adjusted logistic regressions, multiple regressions or ANCOVAs, we extracted the adjusted findings only.

**Table DS4. Medium/long-term effectiveness of interventions to reduce stigma: statistical significance and direction of findings (longitudinal, panel, prospective and cohort studies)**

Study	Country	High or low/middle income country <sup>a</sup>	Design <sup>b</sup>	Participant type	Intervention Participants (n)	Control Participants (n)	Intervention Description	Condition that was focused on	Control description	Time to final follow-up	
Corker 2013	England	HIC	Longitudinal panel study	Mental health service users	T1: 537 T2: 979 T3: 1016	n/a	Time To Change programme which aims to reduce stigma and discrimination against people with mental health problems.	Mental Illness	n/a	3 years	Scale
											Item
Evans-Lacko 2013a	England	HIC	Longitudinal panel study	General population	T1: 1751 T2: 1745 T3: 1717	n/a	Time To Change programme which aims to reduce stigma and discrimination against people with mental health problems.	Mental Illness	n/a	3 years	Scale
											Item
Evans-Lacko 2013b	England	HIC	Longitudinal panel study	General population	T1: 1110 T2: 868	n/a	Social marketing campaign of the Time To Change programme which aims to reduce stigma and discrimination against people with mental health problems.	Mental Illness	n/a	2.5 years	Scale
											Item
Henderson 2012	England	HIC	Longitudinal panel study	Users of specialist mental health services	T1: 537 T2: 1047	n/a	Time To Change programme which aims to reduce stigma and discrimination against people with mental health problems.	Mental Illness	n/a	1 year	Scale
											Item
Henderson 2013	England	HIC	Longitudinal panel study	Senior British employers	T1: 480 T2: 500 T3: 500	n/a	Time To Change programme which aims to reduce stigma and discrimination against people with mental health problems.	Mental Illness	n/a	4 years	Scale
											Item
Crisp 2005	UK	HIC	Longitudinal panel study	General population	T1: no info T2: 1725	n/a	Changing Minds campaign which aimed challenge discrimination and educate the public about mental disorders.	Mental Illness	n/a	5 years	Scale
											Item
Mehta 2009 <sup>d</sup>	UK	HIC	Longitudinal cohort study with control group	General population	T1: 1000 T2: 1000	T1: 1000 T2: 1000	See Me anti-stigma campaign which aimed to deliver specific messages to members of the public using a range of media.	Mental Illness	Negative coverage of mental health legislation issues in England	3 years	Scale
											Item
Paykel 1998	UK	HIC	Longitudinal panel study	General population	T1: 2009 T2: 2050 T3: 1946	n/a	Defeat Depression campaign aimed to reduce stigma associated with depression in order to encourage earlier treatment-seeking using a range of media.	Depression	n/a	6 years	Scale
											Item

Dietrich 2010	Germany	HIC	Longitudinal panel study with control group	General population	T1: 713 T2: 740 T3:709	T1: 710 T2: 750 T3:707	Nuremberg Alliance Against Depression campaign about depression using print materials and local media to counteract negative media reporting.	Depression	Samples from another city in Germany, Wurzberg, where the intervention did not occur	2 years	Scale
					Item						
Gaebel 2008	Germany	HIC	Longitudinal cohort study with control group	General population	T1: 2443 T2: 1493	T1: 2391 T2: 1571	Global Program against Stigma and Discrimination Because of Schizophrenia campaign which promoted contact to people with mental illness and education of target groups.	Schizophrenia	No antistigma or awareness programmes conducted in Berlin and Essen.	3 years	Scale
					Item						
Gaebel 2008 (2nd arm)	Germany	HIC	Longitudinal cohort study with control group	General population	T1: 2391 T2: 1558	T1: 2391 T2: 1571	Awareness programme set up to address specific target groups who are frequently in contact with young people at risk of developing schizophrenia.	Schizophrenia	No antistigma or awareness programmes conducted in Berlin and Essen.	3 years	Scale
					Item						
Grausgruber 2009	Austria	HIC	Longitudinal panel study	General population	T1: 1042 T2: 988	n/a	Anti-stigma campaign with events including: educational interventions in schools, social advertising and conferences where people with mental health problems and family members were able to discuss their experiences.	Mental Illness, focus on Schizophrenia	n/a	9 years	Scale
					Item						
Jorm 2005	Australia	HIC	Longitudinal panel study with control group	General population	T1: 687 T2: 522	T1: 323 T2: 388	Beyondblue campaign which aimed to improve knowledge and responses to depression. Its priority areas include community awareness and destigmatisation.	Depression	Surveys carried out in low-exposure states (those that did not provide funding for the campaign)	8 years	Scale
					Item						
Jorm 2006	Australia	HIC	Longitudinal panel study with control group	General population	T1: 687 T2: 522	T1: 323 T2: 388	Beyondblue campaign which aimed to improve knowledge and responses to depression. Its priority areas include community awareness and destigmatisation.	Depression	Surveys carried out in low-exposure states (those that did not provide funding for the campaign)	8 years	Scale
					Item						
Reavley 2012	Australia	HIC	Longitudinal panel study	General population	T1: 3998 T2: 5860	n/a	Australian national campaigns to improve mental health literacy and reduce stigma: beyondblue and SANE Australia.	Mental Illness	n/a	8 years	Scale
					Item						
Wright 2006	Australia	HIC	Longitudinal panel study	Young people aged 12-25 years	T1: 538 T2: 520	T1: 538 T2: 520	Mental health literacy community awareness campaign targeting young people using multimedia, a website and an information telephone service.	Mental Illness	Regions in which intervention was not implemented	14 months	Scale
				Item							

**a** High income country (HIC) or low to middle income country (LAMIC). Based on World Bank classifications as of July 1, 2012

**b** Study designs included: longitudinal panel studies with a control group, longitudinal panel studies without a control group and longitudinal cohort studies with a control group.

**c** + indicates  $p < 0.05$  in a direction of benefit (reduction in stigma); 0 indicates  $p > 0.05$ ; - indicates  $p < 0.05$  in direction of harm (increase in stigma). Each symbol represents one outcome (scales, or item). Comparisons were made between baseline and final follow-up assessments.

**d** This study found evidence of harm in 4/25 items in the intervention group and 17/25 items in the control group; results are presented as a reduction of harm (benefit) in 13/25 items



Evidence of effectiveness<sup>c</sup>

Knowledge	Attitudes	Behaviour	Outcome measures
		00+	
			The Discrimination and Stigma Scale (Brohan et al., 2012)
0	+	0	Mental Health Knowledge Schedule (Evans-Lacko et al., 2010) Attitudes to Mental Illness questionnaire (Department of Health, 1993 ) Intended Behaviour Scale (Evans-Lacko et al, 2011) Reported Behaviour Scale (Evans-Lacko et al, 2011)
0	0		Mental Health Knowledge Schedule (Evans-Lacko et al., 2010) Attitudes to Mental Illness questionnaire (Department of Health, 1993 ) Intended Behaviour Scale (Evans-Lacko et al, 2011) Reported Behaviour Scale (Evans-Lacko et al, 2011)
	000+		
		+	
			The Discrimination and Stigma Scale (Brohan et al, 2012)
		0++	
000000+++++	00+++++	0000+++++	Knowledge of Mental Health in the Workplace (Henderson et al., 2013) Attitudes of Employers (Henderson et al., 2013) Mental Health Policies in the Workplace; Preparedness for Workplace Mental Health Problems (Henderson et al., 2013)
			Negative Attitudes to People with Mental Illnesses (Crisp et al., 2005)
	00000000000000+++++		
			Community Attitudes to the Mentally Ill survey (Taylor & Dear, 1981)
	0000000000+++++		
			Structured interview (Paykel et al., 1998)
0000+++++	000++++	0+	



**Table D55. Effectiveness of interventions to reduce stigma in low- and middle-income countries: study characteristics, statistical significance and direction of findings**

								Evidence of Effectiveness (at post-intervention) <sup>c</sup>			
Study	Country	Income Group <sup>a</sup>	Design <sup>b</sup>	Participant type	Intervention Participants (n)	Control Participants (n)	Intervention Description		Knowledge	Attitudes	Behaviour
Gutierrez-Maldonado 2009	Chile	Upper-middle income	RCT	Key caregivers of patients with Schizophrenia	18	23	Psychoeducational programme aiming to change caregivers attitudes to schizophrenia and their general health perceptions.	Scale		+	
								Item			
Hao 2011	China	Upper-middle income	RCT	University students majoring in developmental psychology	34	30	Group mental health training for improving students intentions and attitudes towards seeking psychological help.	Scale		++++	
								Item			
Rahman 1998	Pakistan	Lower-middle income	RCT	School children (12-16 years)	50	50	Mental health programme in schools to improve the understanding of mental-health disorders and increase awareness of available treatment. Schoolchildren share knowledge with friends, family and neighbours.	Scale		+	
								Item			
Rahman 1998 (sample 2)	Pakistan	Lower-middle income	RCT	Parents of school children	50	50	As above	Scale		+	
Rahman 1998 (sample 3)	Pakistan	Lower-middle income	RCT	Friends of school children not in school	50	50	As above	Item			
								Scale		+	
Rahman 1998 (sample 4)	Pakistan	Lower-middle income	RCT	Neighbours of school children	50	50	As above	Item			
								Scale		+	
Bayar 2009	Turkey	Upper-middle income	Post-test controlled	Medical professionals in psychiatry	100	105	An instructive e-mail that presented a general account of stigma in mental health.	Scale		+	
								Item			
Altindag 2006	Turkey	Upper-middle income	Controlled	Medical Students	25	35	1 day program with an educational lecture about stigma and schizophrenia, direct contact with a young person with schizophrenia and indirect contact, viewing of the film 'A Beautiful Mind'.	Scale			
								Item		0000+++	
Armstrong 2011	India	Lower-middle income	Uncontrolled	Community health workers	70	n/a	Mental health training program aiming to increase recognition of mental disorders, enhance appropriate response and referral, support people with mental disorders and their families, and improve mental health promotion in communities.	Scale			
								Item	000000000000++++	0	
Chinnayya 1990	India	Lower-middle income	Uncontrolled	Primary health care workers	150	n/a	1 week mental health training programme - lectures, case demonstrations and role play.	Scale			
								Item	0000+++++	0++	
Garcia Silberman 1993	Mexico	Upper-middle income	Uncontrolled	High school students (16-23 years)	21	n/a	A commercial film depicting a true story in which a psychiatrist is portrayed in a positive light.	Scale			
								Item	000000000000000000000000++++		
Garcia Silberman 1993 (2nd arm)	Mexico	Upper-middle income	Uncontrolled	High school students (16-23 years)	21	n/a	A commercial film depicting a true story in which a psychiatrist is portrayed in a negative light.	Scale			
								Item	000000000000++++		
Ghorbani 2009	Iran	Upper-middle income	Uncontrolled	High school students (15-16 years)	639	n/a	A teen health website covering mental health topics in various media formats.	Scale	0		
								Item			
Rong 2011	China	Upper-middle income	Uncontrolled	Medical Students	103	n/a	The intervention consisted in a 1.5 hour lecture which covered all the basic medical aspects about depression required by the teaching guidelines	Scale		0	
								Item		--00000000++	
Rong 2011 (2nd arm)	China	Upper-middle income	Uncontrolled	Medical Students	102	n/a	The intervention consisted in a 1.5 hour lecture about depression and 2 weeks of additional activities carried out by students about the experience of mental illness.	Scale		+	
								Item	000000000++++		
Worakul 2007	Thailand	Upper-middle income	Uncontrolled	Caregivers or family members	91	n/a	Psycho-educational program aiming to provide knowledge, reduce the caregivers' stress, and help	Scale	0		

1998	1998	Upper middle income	Uncontrolled	of patients with schizophrenia	22	1998	the caregivers to provide care to the patients more effectively.	Item		
------	------	---------------------	--------------	--------------------------------	----	------	--	------	--	--

a Categorized using the World Bank classifications as of July 1st 2012

b Type of study design includes: randomised controlled trials (RCT), randomised controlled trials analysed within-group (RCT1), post-test only controlled experimental study (Post-test controlled) and pre-post studies with no control group (Uncontrolled). Findings at post-intervention: comparisons were made between baseline and post-intervention assessments and grouped into knowledge and attitudes.

c + indicates  $p < 0.05$  in a direction of benefit (reduction in stigma); 0 indicates  $p > 0.05$ ; - indicates  $p < 0.05$  in direction of harm (increase in stigma). Each symbol represents one outcome (scales, or item)

Rahman A, Mubbashar MH, Gater R, Goldberg D. Randomised trial of impact of school mental-health programme in rural Rawalpindi, Pakistan. *Lancet*. 1998;352(9133):1022-5.

Bayar MR, Poyraz BC, Aksoy-Poyraz C, Arikan MK. Reducing mental illness stigma in mental health professionals using a web-based approach. *Israel Journal of Psychiatry & Related Sciences*. [Randomized Controlled Trial]. 2009;46(3):226-30.

Premalatha Chinnayya H, Chandrashekar CR, Mooly S, Puttamma, Raghuram A, Subramanya KR, et al. Training Primary Care Health Workers in Mental Health Evaluation of Attitudes towards Mental Illness before and after Training. *International Journal of Social Psychiatry*. 1990 December 1, 1990;36(4):300-7.

Garcia Silberman S. Mental health and the movies: An experimental study about attitude change. *Revista Mexicana de Psicologia*. 1993 Jan-Jun;10(1):37-48.

Ghorbani NR, Heidari RN. Effects of information and communication technology on youth's health knowledge. *Asia-Pacific Journal of Public Health*. 2011 May;23(3):363-8.

Worakul P, Thavichachart N, Lueboonthavatchai P. Effects of psycho-educational program on knowledge and attitude upon schizophrenia of schizophrenic patients' caregivers. *J Med Assoc Thai*. 2007 2007;90(6):199-204.



**Table DS6. Quality ratings for randomised controlled trials using Cochrane Risk of Bias Tool**

Study reference	Random sequence generation	Allocation concealment	Blinding participants and personnel	Blinding of outcome assessment	Incomplete outcome data	Selective outcome reporting
Gulliver 2012	L	L	L	L	L	L
Fung 2011	L	U	U	L	L	L
Jorm 2004*	L	L	H	L	L	L
Jorm 2010a	L	L	H	L	L	L
Kitchener 2004	L	L	H	L	L	L
Aho-Mustonen 2011	L	U	H	L	L	H
Irvine 2012	U	U	U	L	L	L
Jorm 2010b*	L	H	H	L	L	L
Seal 2012	U	U	U	L	L	L
Sharp 2006	U	L	U	L	H	L
Yanos 2012	L	L	U	U	L	L
Kerby 2008	U	L	H	L	L	H
Luty 2008	L	U	U	U	H	L
Pinto-Foltz 2011*	U	U	H	U	L	L
Campbell 2011*	U	U	U	U	H	L
Gumley 2006	L	H	U	H	L	H
Castro 2012*	H	H	H	U	H	U
Gonzales 2002	U	U	U	U	H	L
Link 2002	U	U	U	U	H	L

\*Cluster randomised trials

§Validated = at least one scale was validated; Previously published = scale had been previously published, no evidence of validity

Other bias	Validity of Scales§
U	Validated
L	Validated
U	Validated
U	Validated
U	Validated
L	Validated
L	Validated
H	Validated
L	Previously published
L	Validated
H	Validated
H	Validated
L	Validated
L	Validated
L	Validated
U	Validated
L	Previously published
H	Validated
U	Validated

**Table DS7. Quality ratings for non randomised studies using**

Study reference	Selection Bias (intervention vs control group)¥
Altindag 2006	U
Anderson 2012	n/a
Armstrong 2011	n/a
Buizza 2007	n/a
Conrad 2009	U
Domino 1983	L
Economou 2012	L
Essler 2006	n/a
Esters 1998	U
Faigin 2008	L
Friedrich 2013	H
Gould 2007	H
Graham 2010	n/a
Kelly 2011	n/a
Kitchener 2002	n/a
Knight 2006	U
Krawitz 2004	n/a
Masuda 2009	n/a
Naylor 2009	L
Ng 2002	L
O'Kearney 2006	H
O'Kearney 2009	H
O'Reilly 2010	n/a
O'Reilly 2011	H
Patterson 2007	H
Pejovic-Milovancevic 2009	n/a
Pierce 2010	n/a
Pinfold 2003a	n/a
Pinfold 2003b	n/a
Rong 2011	L

Schneider 2011	n/a	
Schulze 2003		H
Shandley 2010		
Sogaard 1992	n/a	
Treolar 2009		H
Ucock 2006	n/a	
Ventieri 2011		H
Corker 2013	n/a	
Evans-Lacko 2013a	n/a	
Evans-Lacko 2013b	n/a	
Henderson 2012	n/a	
Henderson 2013	n/a	
Crisp 2005	n/a	
Mehta 2009	n/a	
Paykel 1998	n/a	
Dietrich 2010	n/a	
Gaebel 2008		H
Jorm 2005		H
Jorm 2006		H
Reavley 2012	n/a	
Wright 2006		H
Grausgruber 2009		n/a
Yamaguchi 2007		U

¥Comparability between the intervention and control groups. †Repr  
 ΔWhether or not the authors controlled for any confounders in the ;  
 developed by authors with no evidence of validity; Items only= scale

**ing criteria developed by review team (based on an adaptation of the Co**

Selection Bias (representativeness)†	Attrition Bias‡	Selective Outcome reporting
H	L	L
U	H	L
U	L	H
U	H	H
L	U	H
U	H	L
L	L	L
H	L	L
U	L	L
H	L	L
U	H	L
U	H	L
H	H	H
H	H	L
H	L	L
L	L	L
U	H	L
H	L	L
L	L	L
U	H	L
H	L	L
L	H	L
L	H	L
L	L	L
H	H	L
U	L	L
U	H	H
L	H	L
U	H	L
U	U	L

H		L	L
H		U	L
H		L	L
L		H	L
H		H	L
U		H	L
H		L	L
H	n/a		L
L	n/a		L
L	n/a		L
H	n/a		L
L		H	L
H	n/a		L
H	n/a		L
H	n/a		L
L	n/a		L
L		n/a	L
L		L	L

representativeness of sample in comparison to the target population. ‡ Whether there was a factor analysis. § Validated = at least one scale was validated; Previously published = scale has been reported at item level/only single items have been used.



	U	N
	U	Y
	U	N
	U	N
	U	N
	U	N
	U	Y
n/a		N
n/a		Y
n/a		Y
n/a		N
n/a		Y
n/a		N
n/a		Y
n/a		Y
n/a	n/a	Y
	U	N

was more than 20% attrition and if so, whether an intention to treat analysis had been previously published, no evidence of validity; Developed by authc

## Validity of Scales§

Items only

Validated

Items only

Previously published

Validated

Developed by authors

Items only

Previously published

Validated

Validated

Validated

Items only

Validated

Validated

Validated

Validated

Items only

Validated

Developed by authors

Validated

Items only

Validated

Items only

Validated

Validated

Validated

Items only

Items only

Items only

Validated

Items only

Validated

Items only

Items only

Validated

Items only

Validated

Validated

Validated

Validated

Validated

Items only

Items only

Items only

Items only

Developed by authors

Validated

Items only

Items only

Items only

Items only

Developed by authors

Items only

is was carried out.

ors = scale was

# BJP The British Journal of Psychiatry Psych

## **Evidence for effective interventions to reduce mental health-related stigma and discrimination in the medium and long term: systematic review**

N. Mehta, S. Clement, E. Marcus, A.-C. Stona, N. Bezborodovs, S. Evans-Lacko, J. Palacios, M. Docherty, E. Barley, D. Rose, M. Koschorke, R. Shidhaye, C. Henderson and G. Thornicroft  
*BJP* 2015, 207:377-384.

Access the most recent version at DOI: [10.1192/bjp.bp.114.151944](https://doi.org/10.1192/bjp.bp.114.151944)

---

### **Supplementary Material**

Supplementary material can be found at:  
<http://bjp.rcpsych.org/content/suppl/2015/11/09/207.5.377.DC1.html>

### **References**

This article cites 53 articles, 9 of which you can access for free at:  
<http://bjp.rcpsych.org/content/207/5/377#BIBL>

### **Reprints/permissions**

To obtain reprints or permission to reproduce material from this paper, please write to [permissions@rcpsych.ac.uk](mailto:permissions@rcpsych.ac.uk)

### **You can respond to this article at**

[/letters/submit/bjprcpsych;207/5/377](http://bjp.rcpsych.org/letters/submit/bjprcpsych;207/5/377)

### **Downloaded from**

<http://bjp.rcpsych.org/> on January 2, 2016  
Published by [The Royal College of Psychiatrists](http://www.rcpsych.ac.uk)

---